Systematic significance of cypsel morphology in *Lessingianthus* (Vernonieae, Asteraceae)

María B. Angulo^{A,B}, María M. Sosa^A and Massimiliano Dematteis^A

^AInstituto de Botánica del Nordeste (UNNE–CONICET), Casilla de Correo 209, W3400CBL, Corrientes, Argentina.

^BCorresponding author. Email: angulobetiana@gmail.com

Abstract. The taxonomic significance of cypsel features of South American species of *Lessingianthus* (Vernonieae, Asteraceae) is analysed for the first time and discussed in relation to other genera of the tribe Vernonieae. The morphology of the cypselae of 112 species of the genus were analysed using stereo-, light and scanning electron microscopy (SEM) to evaluate the infrageneric relationships and their reliability as taxonomic markers at a generic level. Characters such as cypsel pubescence, carpopodium structure, crystals and idioblasts on the fruit wall were examined. We established three types of cypsel on the basis of the presence or absence, and type of trichomes. Carpopodium is present in all species of the genus. Crystals are very variable in shape and size, with prismatic (rectangular and hexagonal) and styloid shapes. Idioblasts are present in all of the species, except for two. Cypsel features of *Lessingianthus* are often widespread in other related genera of Vernonieae. Therefore, these characters are not good taxonomic markers at the genus level, but they are valuable within genera to differentiate related species from one another.

Additional keywords: Lepidaploinae, taxonomy.

Received 16 May 2015, accepted 7 August 2015, published online 13 November 2015

Introduction

Micromorphological features are a potentially important source of information for classification of the family Asteraceae, being especially useful for the characterisation and classification of tribes and genera (Funk *et al.* 2009). These microcharacters, in combination with other additional characters (type of pollen grains, chromosome number and inflorescence type), were used to classify the tribe Vernonieae. Initially, a subtribal delimitation within Vernonieae was very difficult because most species of tribe were placed in the large genus *Vernonia* (~1000 species). This genus was traditionally defined by the absence of characters present in other Vernonieae genera. In addition, the tribe has a large number of monotypic genera with unusual and distinctive morphological characteristics; therefore, the relationship among those genera and with *Vernonia* was poorly understood. Robinson's (1999) classification changed the circumscription of *Vernonia sens. lat.*, and the genus is currently limited to a small group of 22 taxa from eastern North America, with the other species being in new genera. *Lessingianthus* H. Rob., one of these new genera, arises from the combination of morphological and micromorphological characters.

At present, *Lessingianthus* comprises ~133 species (Angulo and Dematteis 2014) distributed in South America, including Brazil, Bolivia, Paraguay, Argentina, Uruguay, Venezuela, Colombia and Peru (Robinson 2007). Within the tribe, *Lessingianthus* is generally considered to be most closely related to *Chrysolaena* H. Rob. and *Lepidaploa* (Cass.) DC. (Robinson

1999) and molecular studies have supported these relationships (Keeley *et al.* 2007). *Lessingianthus* differs from these genera by a combination of characters, such as 'type B' pollen grains, base chromosome number $x = 16$, and several micromorphological features, such as non-glandular anther apical appendage, absence of basal stylar node (rarely present), and cubic crystals on the fruit wall (Robinson 1999; Angulo and Dematteis 2010, 2012a, 2014). Recently, Angulo and Dematteis (2014) studied in detail the floral microcharacters (corolla pubescence, shape and size of anther apical appendage, style base) of *Lessingianthus* species and demonstrated their usefulness to differentiate related species. However, some features are not diagnostic at generic level because some states of these characters, considered 'typical' of *Lessingianthus*, are also shared with closely related genera such as *Chrysolaena* and *Lepidaploa*. Therefore, the search for new diagnostic characters to help clarify the relationships between these genera is necessary.

In Asteraceae, one of approaches that has shown satisfactory results for taxonomy is the analysis of cypsel morphology (Spjut 1994; Roque and Funk 2013; Zhang *et al.* 2013). The use of cypsel micromorphological characters has proved to be very rewarding for the systematic evaluation of several tribes of Asteraceae, including Anthemideae, Gnaphalieae, Inuleae, Senecioneae, Plucheae and Mutisieae (Abid and Kaiser 2009; Abid and Ali 2010; Abid and Alam 2011; Roque and Funk 2013; Zhang *et al.* 2013). In Vernonieae, cypsel features vary with the group. For example, only the subtribe Sipolisiinae has cypsel

walls containing phytomelanin, a characteristic that is absent in other subtribes (Funk *et al.* 2009). The presence or absence and shape of crystals on the cypsel wall were useful to separate related genera, such as *Vernonanthura* H. Rob. and *Lepidaploa* (Funk *et al.* 2009). The persistence of the pappus is also important at a generic level within the subtribes, such as within the tribe Lychnophorinae, where the genus *Lychnophora* Mart. has a deciduous pappus, whereas *Minasia* H. Rob. has a persistent pappus. The carpopodium is the scar of the abscission zone on the fruit base, and this is another diagnostic character used at different taxonomic levels, showing differences in the size and shape of wall cells (Funk *et al.* 2009). Studies at the infrageneric level are scarce; however, fruit morphology has been used in taxonomic keys to differentiate related species (Angulo and Dematteis 2012b; Bunwong *et al.* 2014; Lander and Hurter 2013). Despite the high number of *Lessingianthus* species, only one study on the morphology and anatomy of the cypselae was conducted (Martins and Oliveira 2007). These authors compared the fruit morphology and anatomy of *L. brevifolius* (Less.) H. Rob. and *Chrysolaena obovata* (Less.) Dematt. (sub nom. *C. herbacea* (Vell.) H. Rob.) and demonstrated the significance of these features to differentiate related taxa. However, there is still no detailed report available on cypsel micromorphology for *Lessingianthus*. In the present paper, the fruits of almost all the species of the genus were analysed in a taxonomic context. Our results will contribute to the characterisation of the species, establishing the value of fruit characters as diagnostic of *Lessingianthus*. In addition, these data are also discussed in relation to the information available on the closely related genera *Chrysolaena* and *Lepidaploa*.

Materials and methods

Cypsel morphology of a total of 112 species was studied with stereo-, light and scanning electron microscopy (SEM). At least five different cypselae per specimen were examined. In total, three specimens per species were analysed depending on material availability. Mature cypselae were collected from plants in the wild or from herbarium specimens (i.e. voucher specimens) from the herbaria ALCB, BR, CESJ, CTES, ESA, G, LP, MBM, P, R, RB, SPF, SI and UB (herbarium abbreviations are according to *Index Herbariorum*, see <http://sweetgum.nybg.org/ih>, accessed 20 April 2015). Species and specimen information is listed in Appendix 1, 'Specimens analysed', where the taxa are arranged alphabetically (Table 1).

For observation of cypsel microcharacters, the fruit was softened in distilled water for 5 min, and cleared in sodium hypochlorite: water (1 : 1) until the samples became transparent. Transparent samples were washed in running water and dissected under a stereomicroscope and then mounted in Hoyer's solution (Anderson 1954; King and Robinson 1970). Samples were examined and photographed using Zeiss Axioplan microscope (Carl Zeiss, Jena, Germany) with digital camera Canon Power Shot A 640 (Tokyo, Japan).

For the anatomical studies, fruit samples were dehydrated following the methodology by Gonzalez and Cristóbal (1997), and were subsequently infiltrated in paraffin (Johansen 1940). Transverse and longitudinal sections were cut into 10–15- μ m sections with a rotary microtome and mounted on glass slides.

Sections were stained with a safranin–astra blue combination (Luque *et al.* 1996) and photographed using a Leica DM LB2 light microscope (Leica Microsystems, Wetzlar, Germany) including polarised light.

For histochemical analyses, living material was hand-sectioned under a stereomicroscope. Cellular content was tested histochemically using Sudan IV for lipids, Lugol's reagent for starch grains, Ruthenium red for pectinaceous materials, and cresyl blue for mucilages. Sections were observed and photographed using light microscope.

The chemical identity of the crystals was tested by the acid treatment described by Zarlavsky (2014). Fruits were softened in distilled water for 5 min and dissected under a stereomicroscope, and a few drops of 2% hydrochloric acid were added.

The different types of crystals were measured, considering length and diameter. One hundred crystals for preparations and each crystal type were measured from 10 randomly selected regions.

For observation of cypsel surface sculpturing, mature fruits were first dehydrated and immersed in CO₂ for critical-point drying before sputter-coating with gold–palladium. Coated samples were examined and photographed using a JEOL 5800 LV scanning electron microscope (JEOL Ltd, Tokyo, Japan).

The term cypsel was used for *Lessingianthus* fruits on the basis of the terminology suggested by Marzinek *et al.* (2008). Terminology of trichome morphology follows Metcalfe and Chalk (1979), whereas terminology of remaining cypsel characters follows Robinson (1999), and Roque *et al.* (2009).

The Infostat software package (Di Rienzo *et al.* 2013) was used for comparative analysis of cypsel characters of *Lessingianthus* species. A principal coordinate analysis (PCoA) using Manhattan distance metric was performed to evaluate the variability of fruit characters among examined species. A data matrix of 112 operational taxonomic units (OTUs) and four qualitative variables was constructed (data matrix is available on request from the senior author). The different character states for each of them are given in Table 2.

Results

Cypselae of *Lessingianthus* vary between cylindrical and obconical, and some species show an intermediate shape designated here as cylindrical–obconical (see Table 1). However, fruit length (without pappus) is highly variable, ranging from 1.5 to 9 mm. *Lessingianthus exiguus* has the smallest fruit length (1.5–1.8 mm), whereas *L. monocephalus* has the largest value (8–9 mm). All the species have more or less conspicuous longitudinal ribs. The pappus is whitish or yellowish, biseriate, with an outer series of short paleaceous setae (0.5–2 mm), fimbriate, lanceolate (Fig. 1A) to linear, and an inner series with long cylindrical bristles (4.5–12 mm).

Trichomes

Cypsel surface in most of the analysed species shows trichomes, although some taxa have glabrous fruits (Table 1). The trichomes are multicellular, glandular or non-glandular. On the basis of the presence of trichomes as well as the observed types, we classify three fruit types, as defined below.

Table 1. Characteristics of cypselas in *Lessingianthus* species

Fruit types: 1, glabrous cypselas; 2, cypselas with twin hairs; 3, cypselas with twin hairs and glandular trichomes. Idioblasts: 1, present; 2, absent. Crystals: 1, rectangular prismatic crystals; 2, hexagonal prismatic crystals; 3, styloid crystals

Species	Fruit shape	Fruit length (mm) (without pappus)	Fruit types			Idioblasts			Carpopodium (average) maximum			Crystals			Species code
			1	2	3	1	2	3	Length (mm)	Width (mm)	3	1	2	3	
<i>L. adenophyllus</i>	Cylindrical	2.5–3		x		x			0.1 (0.15) 0.2	0.18 (0.20) 0.25		x			cf g
<i>L. ammophyllus</i>	Obconical	3–3.5		x			x		0.15 (0.2) 0.25	0.37 (0.40) 0.45		x			co o
<i>L. arachniotepis</i>	Obconical	3–4		x		x			0.15 (0.2) 0.25	0.35 (0.40) 0.45			x		cce
<i>L. arctatus</i>	Obconical	2–2.5		x		x			0.1 (0.15) 0.2	0.33 (0.40) 0.45		x			cte
<i>L. argenteus</i>	Obconical	3.5–4		x		x			0.1 (0.15) 0.2	0.45 (0.50) 0.55		x			arge
<i>L. argyrophyllus</i>	Obconical	3–3.5		x		x			0.1 (0.15) 0.2	0.42 (0.45) 0.50			x	x	argy
<i>L. asteriflorus</i>	Obconical	3.5–4.5	x			x			0.2 (0.25) 0.3	0.48 (0.55) 0.60		x			ast
<i>L. bardanoides</i>	Cylindrical–obconical	4–5		x		x			0.2 (0.3) 0.35	0.59 (0.60) 0.65		x			bard
<i>L. barrosoanus</i>	Obconical	4–4.5	x			x			0.1 (0.125) 0.15	0.50 (0.55) 0.58		x			barr
<i>L. bakerianus</i>	Cylindrical	3–3.3			x	x			0.15 (0.2) 0.2	0.29 (0.30) 0.31		x			bak
<i>L. bellulus</i>	Cylindrical–lobconical	7–8		x		x			0.15 (0.2) 0.25	0.25 (0.30) 0.34		x			bel
<i>L. bishopi</i>	Cylindrical	3–3.5		x		x			0.1 (0.15) 0.2	0.29 (0.30) 0.34		x			bis
<i>L. brevifolius</i>	Cylindrical	3–3.4			x	x			0.1 (0.2) 0.25	0.20 (0.25) 0.31		x			bre
<i>L. brevipetiolatus</i>	Cylindrical	3–3.5		x		x			0.1 (0.15) 0.2	0.4 (0.45) 0.5		x			brevip
<i>L. buddleifolius</i>	Cylindrical–obconical	3–3.5		x		x			0.1 (0.15) 0.2	0.51 (0.55) 0.57		x			bud
<i>L. cardioides</i>	Cylindrical–obconical	3–3.5		x		x			0.1 (0.2) 0.2	0.41 (0.45) 0.46		x			card
<i>L. carvalhoi</i>	Obconical	2–2.3			x	x			0.1 (0.15) 0.2	0.39 (0.40) 0.45		x			carv
<i>L. cataractum</i>	Cylindrical	3–3.5		x		x			0.25 (0.3) 0.4	0.40 (0.45) 0.46		x			cata
<i>L. centaurosideus</i>	Cylindrical–obconical	3–4			x	x			0.1 (0.15) 0.2	0.11 (0.15) 0.20		x			cent
<i>L. cephalotes</i>	Obconical	2–2.4		x		x			0.1 (0.15) 0.2	0.32 (0.35) 0.39		x			cep
<i>L. clavatus</i>	Obconical	2.5–3		x		x			0.1 (0.15) 0.2	0.25 (0.3) 0.32		x			cla
<i>L. compactiflorus</i>	Obconical	2.5–4.3		x		x			0.4 (0.55) 0.6	0.55 (0.6) 0.65		x			com
<i>L. conscriptus</i>	Cylindrical–obconical	4–5		x		x			0.1 (0.15) 0.15	0.29 (0.35) 0.37		x			con
<i>L. cordiger</i>	Obconical	2.5–3		x		x			0.2 (0.25) 0.25	0.26 (0.3) 0.31		x			cor
<i>L. coriaceus</i>	Cylindrical–obconical	3–3.5	x			x			0.1 (0.15) 0.2	0.41 (0.45) 0.48		x			coriac
<i>L. coriarius</i>	Cylindrical–obconical	3–3.5				x			0.14 (0.15) 0.2	0.26 (0.3) 0.34		x			coriar
<i>L. correntinus</i>	Cylindrical–obconical	4–5	x			x			0.1 (0.15) 0.2	0.7 (0.75) 0.8		x			cor
<i>L. cristallinae</i>	Obconical	2.5–3.3		x		x			0.14 (0.15) 0.21	0.5 (0.55) 0.56		x			cri
<i>L. durus</i>	Obconical	3–4		x		x			0.09 (0.1) 0.14	0.28 (0.30) 0.35		x			dur
<i>L. elegans</i>	Obconical	2–2.2		x		x		x	0.07 (0.075) 0.1	0.09 (0.10) 0.11		x			ele
<i>L. exiguus</i>	Cylindrical–obconical	1.5–1.8		x		x			0.05 (0.05) 0.1	0.36 (0.4) 0.5		x			exi
<i>L. farinosus</i>	Obconical	2.5–3		x		x			0.1 (0.2) 0.3	0.30 (0.35) 0.38		x			far
<i>L. floccosus</i>	Cylindrical	3–3.2		x		x			0.1 (0.15) 0.2	0.30 (0.4) 0.45		x			flo
<i>L. foliosus</i>	Obconical	2–2.5		x		x			0.04 (0.05) 0.09	0.21 (0.25) 0.28		x			fol
<i>L. glabratus</i>	Obconical	4–5		x		x			0.1 (0.15) 0.2	0.35 (0.4) 0.45		x			glab
<i>L. grandiflorus</i>	Obconical	4–4.5		x		x			0.02 (0.025) 0.05	0.25 (0.3) 0.35		x			gran
<i>L. grearii</i>	Cylindrical	3.5		x		x			0.12 (0.125) 0.15	0.35 (0.4) 0.41		x		x	gre
<i>L. hasslerianus</i>	Obconical	1.8–2		x		x			0.07 (0.075) 0.10	0.2 (0.25) 0.3		x			has
<i>L. hoveaefolius</i>	Obconical	3–3.3		x		x			0.1 (0.15) 0.2	0.41 (0.45) 0.49		x			hov
<i>L. hypochaeris</i>	Obconical	2		x		x			0.1 (0.15) 0.2	0.16 (0.2) 0.25		x			hyp
<i>L. hystriococcus</i>	Obconical	3–3.5			x	x			0.1 (0.2) 0.3	0.39 (0.4) 0.5		x			hys
<i>L. ibitipocensis</i>	Cylindrical	4–5			x	x			0.2 (0.25) 0.3	0.28 (0.3) 0.35		x			ibi
<i>L. intermedium</i>	Cylindrical	2.5–3		x		x			0.1 (0.2) 0.25	0.40 (0.45) 0.46		x			int
<i>L. irwinii</i>	Cylindrical–obconical	2.5–3		x		x			0.1 (0.15) 0.2	0.5 (0.55) 0.6		x			irw
<i>L. ixiamensis</i>	Obconical	2.5–3		x		x			0.05 (0.1) 0.2	0.34 (0.35) 0.36		x			ixi
<i>L. kuntzei</i>	Cylindrical–obconical	3		x		x			0.2 (0.25) 0.3	0.6 (0.65) 0.70		x			kun

(continued next page)

Table 1. (continued)

Species	Fruit shape	Fruit length (mm) (without pappus)	Fruit types			Idioblasts			Carpopodium (minimum (average) maximum) Length (mm) Width (mm)			Crystals			Species code
			1	2	3	1	2	3	1	2	3	1	2	3	
<i>L. laevigatus</i>	Obconical	1.8–2		x		x			0.1 (0.15) 0.2	0.19 (0.2) 0.25		x			lae
<i>L. lanatus</i>	Obconical	3–3.3		x		x			0.1 (0.15) 0.2	0.29 (0.30) 0.34		x			lana
<i>L. laniferus</i>	Obconical	2.5–3			x				0.1 (0.15) 0.2	0–25 (0.30) 0.37			x		lani
<i>L. lanuginosus</i>	Obconical	3–3.5		x		x			0.05 (0.1) 0.15	0.4 (0.45) 0.48		x			lanu
<i>L. lapinhensis</i>	Obconical	2–2.5		x		x			0.1 (0.125) 0.15	0.2 (0.25) 0.3		x			lap
<i>L. laurifolius</i>	Obconical	3–3.5		x		x			0.1 (0.15) 0.2	0.3 (0.4) 0.45		x			lau
<i>L. ligulaefolius</i>	Obconical	3.5–4		x		x			0.2 (0.25) 0.3	0.8 (0.85) 0.9		x			lig
<i>L. linearifolius</i>	Obconical	2–2.5		x		x			0.1 (0.15) 0.2	0.4 (0.45) 0.5		x			linearif
<i>L. linearis</i>	Obconical	2.5–3		x		x			0.2 (0.25) 0.3	0.2 (0.3) 0.4		x			linear
<i>L. longicauspis</i>	Cylindrical	3.5–4		x		x			0.1 (0.2) 0.2	0.3 (0.4) 0.45		x			lon
<i>L. lorentzii</i>	Cylindrical–obconical	3–4		x		x			0.21 (0.25) 0.3	0.30 (0.35) 0.37		x			lor
<i>L. macrocephalus</i>	Cylindrical	5–6			x				0.1 (0.15) 0.2	0.7 (0.75) 0.8		x			macro
<i>L. macrophyllus</i>	Obconical	4–5		x		x			0.2 (0.25) 0.3	0.4 (0.45) 0.55			x		macro
<i>L. mansoanus</i>	Obconical	5–6		x		x			0.1 (0.2) 0.25	0.4 (0.5) 0.55		x			man
<i>L. membranifolius</i>	Cylindrical–obconical	2.5–3		x		x			0.15 (0.2) 0.25	0.3 (0.35) 0.4		x			mem
<i>L. minimus</i>	Cylindrical–obconical	2.5–3		x		x			0.2 (0.25) 0.3	0.45 (0.5) 0.5		x			min
<i>L. mollissimus</i>	Cylindrical	3–4		x		x			0.2 (0.25) 0.3	0.5 (0.55) 0.6		x			mol
<i>L. monocephalus</i>	Cylindrical	8–9		x		x			0.2 (0.25) 0.3	0.63 (0.65) 0.66		x			mon
<i>L. morii</i>	Cylindrical–obconical	2.5–3	x			x			0.2 (0.25) 0.3	0.2 (0.3) 0.4		x			mor
<i>L. myrsinites</i>	Obconical	4–4.5		x		x			0.2 (0.25) 0.3	0.25 (0.3) 0.4		x			myr
<i>L. niederleinii</i>	Obconical	3.5–4		x		x			0.05 (0.075) 0.1	0.2 (0.3) 0.35		x			nie
<i>L. obscares</i>	Cylindrical–obconical	2.5–2.8		x		x			0.1 (0.2) 0.25	0.2 (0.25) 0.3		x			obs
<i>L. obtusatus</i>	Obconical	2.5–3		x		x			0.08 (0.1) 0.15	0.3 (0.35) 0.4		x			obt
<i>L. onopordoides</i>	Cylindrical	4–4.5		x		x			0.3 (0.35) 0.45	0.5 (0.55) 0.6		x			ono
<i>L. parvifolius</i>	Obconical	2.5		x		x			0.29 (0.30) 0.32	0.3 (0.38) 0.4			x		par
<i>L. pentacanthus</i>	Obconical	3–3.5			x				0.2 (0.25) 0.3	0.7 (0.75) 0.8		x			pen
<i>L. plantaginoides</i>	Cylindrical–obconical	2.5–3		x		x			0.2 (0.30) 0.35	0.4 (0.45) 0.5		x			plan
<i>L. platyphyllus</i>	Obconical	3–3.5		x		x			0.07 (0.075) 0.1	0.15 (0.2) 0.3		x			plat
<i>L. polyphyllus</i>	Obconical	2–2.5		x		x			0.1 (0.15) 0.2	0.2 (0.30) 0.4		x			pol
<i>L. profusus</i>	Cylindrical–obconical	4–4.5		x		x			0.07 (0.075) 0.1	0.2 (0.3) 0.4		x			pro
<i>L. pseudoincanus</i>	Obconical	3–3.5			x				0.1 (0.15) 0.2	0.2 (0.25) 0.3		x			pse
<i>L. psilophyllus</i>	Obconical	3–4		x		x			0.1 (0.2) 0.3	0.2 (0.25) 0.3		x			psi
<i>L. pubescens</i>	Obconical	2.8–3.2		x		x			0.15 (0.20) 0.21	0.3 (0.35) 0.4		x			pub
<i>L. pumilus</i>	Cylindrical	5–8		x		x			0.2 (0.3) 0.4	0.3 (0.35) 0.4		x			pum
<i>L. pusillus</i>	Cylindrical–obconical	3–3.6		x		x			0.1 (0.125) 0.2	0.2 (0.3) 0.4		x			pus
<i>L. pycnostachyus</i>	Cylindrical	4.5–5		x		x			0.1 (0.125) 0.2	0.2 (0.25) 0.3		x			pyc
<i>L. ramellae</i>	Cylindrical–obconical	4–4.5		x		x			0.1 (0.125) 0.2	0.2 (0.25) 0.3		x			ram
<i>L. regis</i>	Obconical	2		x		x			0.05 (0.075) 0.1	0.2 (0.25) 0.3		x			reg
<i>L. reitzianus</i>	Cylindrical–obconical	3–3.5		x		x			0.2 (0.25) 0.3	0.3 (0.35) 0.4		x			rei
<i>L. robustus</i>	Obconical	3–4		x		x			0.1 (0.2) 0.3	0.45 (0.5) 0.6		x			rob
<i>L. roseus</i>	Cylindrical–obconical	2–2.5		x		x			0.2 (0.25) 0.3	0.5 (0.6) 0.7		x			ros
<i>L. rosmarinifolius</i>	Obconical	2–2.5			x				0.09 (0.1) 0.2	0.3 (0.35) 0.4			x		rosm
<i>L. rubraulis</i>	Obconical	3–3.5			x				0.1 (0.15) 0.2	0.25 (0.3) 0.35		x			rub
<i>L. saltenis</i>	Obconical	2.5–3		x		x			0.25 (0.3) 0.4	0.35 (0.4) 0.5		x			sal
<i>L. sancti-pauli</i>	Obconical	2.5–3		x		x			0.9 (0.1) 0.15	0.25 (0.3) 0.4			x		san
<i>L. samosti</i>	Obconical	3–3.5		x		x			0.1 (0.2) 0.25	0.45 (0.5) 0.51		x			sant
<i>L. scabrifolius</i>	Cylindrical–obconical	3–3.5			x				0.15 (0.2) 0.3	0.4 (0.45) 0.5		x			sea
<i>L. secundus</i>	Obconical	2–2.5		x		x			0.05 (0.075) 0.1	0.1 (0.2) 0.25		x			sec
<i>L. sellowii</i>	Cylindrical–obconical	2.4–2.6		x		x			0.02 (0.05) 0.1	0.26 (0.30) 0.35			x		sel

Table 2. List of characters and their coding states

Character number	Fruit character
1	Fruit shape: cylindrical (1), cylindric-obconical (2), obconical (3)
2	Fruit types: glabrous cypselae (1), cypselae with twin hairs (2), cypselas with twin hairs and glandular trichomes (3)
3	Idioblasts: absent (0), present (1)
4	Crystals: rectangular prismatic (1), rectangular and hexagonal prismatic (2), hexagonal prismatic and styloid (3)

Fig. 1. Cypselae microcharacters of *Lessingianthus* (scanning electron micrographs): A, B. Glabrous cypselae of *L. varroniifolius*. A. Glabrous fruit with the external series of pappus (arrow). B. Fruit detail, showing the idioblasts and ribs. C. Detail of carpodium, showing twin hairs (arrow) of *L. bardanoides*. D–F. Cypselae of *L. ulei*. D. Detail of fruit base, showing twin hairs and glandular (arrow) trichomes. E. glandular trichome. F. Detail of carpodium, showing subquadrate cells. G, H. Glabrous cypselae of *L. coriarius*. G. Cypselae base showing carpodium and idioblasts. H. Idioblasts scattered and longitudinally arranged along the sides of the ribs, forming groups of up to nine idioblasts. I. Idioblasts of *L. varroniifolius*.

Fig. 2. Cypsela microcharacters of *Lessingianthus* (light micrographs). A, B. Twin hairs of *L. bardanoides*. C. Twin hairs of *L. argyrophyllus*. D, E. Cypsela of *L. macrocephalus*. D. Detail of the fruit, showing glandular trichomes (arrow) on the carpodium apex. E. Glandular trichomes on the fruit. F. Fruit detail of *L. brevifolius*, showing glandular trichomes (arrow) on the carpodium apex. G. Detail of fruit of *L. grearii*, showing twin hairs and idioblasts. H. Idioblasts (in groups of two and three) of *L. laniferus*. I. Solitary idioblasts of *L. ulei*. Scale bars: 50 μ m (A–C, E, H, I), 100 μ m (D, F, G).

Idioblasts are larger than the cells of the uniseriate epidermis, emerging clearly from the fruit wall (Figs 1H, I, 2G–I). In some species (such as *L. myrsinoides*), they can be distinguished under stereomicroscope. They are scattered (Fig. 2I) or longitudinally arranged along the sides of the ribs, forming groups of up to nine idioblasts in *L. coriarius* (Fig. 1H).

The contents of the idioblasts were submitted to histochemical tests with Lugol's reagent, Sudan IV, Ruthenium red and cresyl

blue with negative results, indicating the absence of starch grains, lipids, pectinaceous materials and mucilages respectively.

Statistical analysis

Principal coordinate analysis (PCoA) of the cypsela characters showed that the first two principal coordinates represent only 45.6% of the total variation (see Fig. 5 for a bidimensional

Fig. 3. Crystals of fruit wall of *Lessingianthus* (light micrographs). A. Rectangular prismatic crystals of *L. durus*. B. Detail of fruit, showing rectangular prismatic crystals of *L. laniferus*. C. Rectangular prismatic crystals (short and long) of *L. glabratus*. D. Short rectangular prismatic crystals of *L. syncephalus*. E, F. Hexagonal prismatic crystals of *L. exiguus*. G. Hexagonal (empty arrow) and rectangular prismatic (filled arrows) crystals of *L. plantaginodes*. H. Styloid crystals (arrow) of *L. argyrophyllus*. I. Styloid crystals of *L. grearii*. Scale bars: 50 μm (A, C, D–I), 100 μm (B).

Fig. 4. Transversal sections of fruits (light micrographs). A. Detail of the rib, showing idioblast (arrow) in the epidermis of *L. rubicaulis*. B. Detail of the idioblasts (arrows) in the epidermis of *L. saltensis*. Scale bar: 50 µm.

Fig. 5. Graph of the first two principal coordinates of *Lessingianthus* species on the basis of cypselal characters. Circles show the grouped species by fruit characters and several subgroups (Subgroups 1A–G, 2A–C) are distinguished within. Subgroups include species with the same coordinates (see text). Codes identifying species are in Table 1.

projection of the axes). However, two distinct groups were differentiated. The large Group 1 includes 76 species (67.85%), which have rectangular prismatic crystals and idioblasts. Several subgroups containing species with the same distribution (same

coordinates) were distinguished within. The same coordinates for species indicate that they share the same micromorphological characteristics. All taxa of the Subgroups 1A (with 31 species), 1B (with 12 species) and 1C (with 14 species) also have cypselae

with twin hairs but differ in the shape of the fruit. Subgroup 1A has obconical cypselae, whereas Subgroup 1B has cylindrical–obconical fruits and Subgroup 1C has cylindrical cypselae. Subgroups 1D (5 species), 1E (3 species) and 1F (4 species) show cypselae with twin hairs and glandular trichomes and also differ in fruit shape, as follows: obconical (Subgroup 1D), cylindrical–obconical (Subgroup 1E) and cylindrical (Subgroup 1F). The only species with glabrous cylindrical cypselae were grouped in Subgroup 1G (5 species). Two species (*L. barrosoanus* and *L. asteriflorus*) were separated from these subgroups because they had a combination of characters, such as obconical and glabrous fruits.

Group 2 is smaller (with 32 species) than Group 1 and it includes taxa with cypselae with idioblasts and rectangular and hexagonal prismatic crystals. This group also has several subgroups within it; Subgroups 2A (12 species) and 2B (10 species) both have cypselae with twin hairs, but fruits are obconical in 2A and cylindrical–obconical in 2B. Subgroup 2C, with five species, has obconical cypselae with twin hairs and also glandular trichomes. Several taxa have a combination of characters, which does not correspond to the characteristics of the subgroups above. For example, *L. centaurospideus* shares some characteristic with Subgroup 2B but it has fruits with twin hairs and glandular trichomes. *Lessingianthus cataractarum* and *L. intermedius* were separated from the rest of the species because they had cylindrical cypselae.

Lessingianthus syncephalus and *L. argyrophyllus* were also isolated by having hexagonal prismatic and styloid crystals.

Lessingianthus grearii was separated (on the negative extreme of coordinate 1) from most of species, because it has cylindrical fruits with crystals hexagonal in shape, prismatic and styloids.

Three species were isolated from the main groups (on the positive extreme of coordinate 1) by lacking idioblasts. *Lessingianthus durus* and *L. ammophyllus* have obconical fruits with rectangular prismatic crystals but differed in the trichome type on fruits, twin hairs and glandular trichomes in the former and only twin hairs in the latter. In contrast, *Lessingianthus soderstroemii* has cylindrical–obconical fruits with twin hairs and rectangular prismatic crystals.

Discussion

Several studies have emphasised the importance of cypselae features for systematic investigation in Asteraceae, because they are useful to clarify the delimitation and establish the similarity of taxa in the family (Grau 1977; Robinson and King 1977; Anderberg 1989; Bremer 1994). Cypselae features, including the presence and number of the ribs, crystals in epidermal cells, carpodium structure, pappus type and trichomes, are used in these classifications because they are stable and genetically controlled (Ciccarelli *et al.* 2007). In the present paper, we studied 112 species of *Lessingianthus* to describe and explore whether cypselae features have diagnostic value for taxonomical purposes.

Lessingianthus species show similar characteristics in general morphology of cypselae to those of other genera of Vernoniaceae (Robinson 1999; Funk *et al.* 2009). For example, twin hairs on fruit of *Lessingianthus* were also observed in all Vernoniaceae, being considered the trichome type that characterises Asteraceae cypselae. Twin hairs, also called double hairs and

‘Zwillingshaare’ (Kraus 1866), have been extensively studied in the family. Their morphology shows great variation in size, length, wall thickness, degree of divergence and degree of development of the cells and the basal cells (Freire and Katinas 1995; Sancho and Katinas 2002; Ike and Nordenstam 2012). In *Lessingianthus*, the morphology of twin hairs varies in length and the degree of divergence of parallel cells; however, the overall morphology fits the description proposed by Cabrera (1944) for *Vernonia* twin hairs. Twin hairs have also been reported in corollas of some species of Asteraceae, such as *Encelia* Adans. (Heliantheae; Carpenter 1999) and in several genera of Mutisieae (Sancho and Katinas 2002). However, this trichome type was not observed in corollas of *Lessingianthus* (Angulo and Dematteis 2014) and is restricted only to the fruits.

The genus was found to have three fruit types on the basis of the presence or absence of trichomes, and the trichome type, with the cypselae with twin hairs being the most common type. Glabrous fruits were observed in eight species, whereas the presence of glandular trichomes, along with non-glandular trichomes, was found in 19 species. In Asteraceae, glandular trichomes of the fruit are usually formed by a stalk and a head of one or many cells (Narayana 1979). Glandular trichomes of *Lessingianthus* share this general structure with a 3–5-celled uniseriate stalk and a unicellular head. A biseriate 10-celled glandular trichome is considered the most common type in Asteraceae and occurs in the tribes Anthemideae (Werker *et al.* 1994), Astereae (Castro *et al.* 1997), Eupatorieae (Castro *et al.* 1997), Heliantheae (Castro *et al.* 1997), Inuleae (Werker and Fahn 1981), Mutisieae (Castro *et al.* 1997) and Vernoniaceae (Faust and Jones 1973; Narayana 1979; Castro *et al.* 1997). However, this glandular trichome type was not found in *Lessingianthus* fruits. The structure of glandular trichomes here described agrees with one type (designated as Type III) on corollas of some species of the genus (Angulo and Dematteis 2014). The secretion of lipids by the trichomes of corollas was observed via histochemical tests (Angulo and Dematteis 2014). However, our findings indicated that glandular trichomes of cypselae do not secrete lipids, nor starch, mucilage or pectinaceous materials. In general, the secretions may have many functions, such as attracting pollinating insects to flowers, and will protect the plant and its reproductive organs against pathogens and herbivores (Ciccarelli *et al.* 2007). Therefore, a detailed study of secretory material, with observations of plants in their natural habitat and chemical analysis, can provide a sound basis for speculation about the true function of these fruit trichomes.

The carpodium may show taxonomically useful variation in size and shape of cells and thickness of cell walls (King and Robinson 1970). However, in *Lessingianthus*, carpodia show a comparative uniformity; all the species have this structure as a complete ring without interruptions, although differences in size (length and width) of this ring were observed. Our results agree with the annular carpodium cited for some species of *Vernonia* (Haque and Godward 1984), and other genera of Vernoniaceae (Funk *et al.* 2009). Within the tribe, only the genus *Camchaya* Gagnep. has no evident carpodium (Funk *et al.* 2009). In other tribes, the carpodium exhibits greater variability, for example in *Wedelia* Jacq. (Heliantheae), the shape may change as the fruit matures, and in Ayapaninae (Eupatorieae), the basal row of cells is greatly enlarged (Funk *et al.* 2009). Several functions have

been assigned to the carpopodium and all of them are related to seed dispersal. Haque and Godward (1984) suggested that the thickened cells provide a rigid structure which, on drying, will not contract, resulting in tension to the thin-walled cells of the abscission layer, making abscission easier. The function of the carpopodium in *Lessingianthus* is still unknown, but we consider that its presence is also related to seed dispersal.

The presence of CaOx crystals in the cypsel wall was observed in all *Lessingianthus* species. These crystals are common for most vegetative and reproductive organs in plants and are widely distributed in over 215 plant families (Franceschi and Horner 1980; Molano-Flores 2001). However, there are only a few studies related to their presence in Asteraceae (Dormer 1961; Horner 1977; Meric 2008; Meric and Dane 2004). In the present study, prismatic (rectangular and hexagonal) and styloid crystals were observed in *Lessingianthus* species, with the prismatic shape being the most frequent. Robinson (1988a) established that the genus is characterised by quadrate crystals (the correct term would be cubic because it is a tridimensional structure), whereas *Lepidaploa* and *Chrysolaena* have elongated crystals (the correct term would be prismatic). Therefore, the crystal shape is one of the diagnostic features that distinguishes *Lessingianthus* from those related genera (Robinson 1988a, 1988b). However, we observed a wide variation in the shape and size of the crystals within the genus. Prismatic crystals were the most common, mainly the rectangular crystals, whereas hexagonal shapes were observed in a few species. Only two species had styloid crystals. Therefore, the crystal shape of cypselae is not a good diagnostic character for *Lessingianthus* because it is very variable and overlapping in most Vernoniae genera.

The mechanisms controlling the morphology of crystals are unknown (Nakata 2003). The diversity of crystal shapes, as well as their prevalence and spatial distribution, has led to several hypotheses about their functions in plants. Some of the proposed functions include roles in ion balance, plant defence, detoxification and tissue support (Franceschi and Horner 1980).

The presence of crystals in both vegetative and reproductive organs was observed in various taxa of Asteraceae (Dormer 1961; Meric and Dane 2004; Meric 2008). It is probable that other reproductive (anthers, petals) and vegetative (leaves, stems, and roots) structures also have crystals in *Lessingianthus*. Therefore, further studies that include other reproductive and vegetative organs are necessary in the genus, so as to determine their presence and, consequently, their morphology and distribution.

There are studies showing that the process of CaOx crystal formation may occur in specialised cells, called crystal idioblasts (Mazen *et al.* 2004). Although the fruit wall of Vernoniae often has idioblasts, there are no studies linking these cells with production of crystals. Almost all of *Lessingianthus* species showed idioblasts in the cypsel wall, which were clearly distinguishable from trichomes and epidermal cells. *Lessingianthus durus* and *L. soderstroemii* had cypselae without idioblasts; however, both species have crystals. Therefore, the observed idioblasts would not be involved in crystal formation. Also, histochemical tests performed here showed that idioblasts do not contain starch grains, lipids, pectinaceous materials or mucilage.

These cells differ from the surrounding tissue cells by their shape, size and wall structure. Hind (1993) described such cells as

glands; however, our results indicated that these cells are certainly idioblasts as suggested by Robinson (1999). Such cells differ from the structure of glandular trichomes (uniserial stalk and unicellular head) observed in *Lessingianthus* fruits. Their presence or absence has been used as a taxonomic feature for Vernoniae (Isawumi *et al.* 1996; Robinson 2009). Almost all species of *Lessingianthus* have idioblasts (only two species do not have these cells); therefore, they are a good taxonomic marker at an infrageneric level.

Taxonomy

Our findings suggested that cypsel features are useful in distinguishing species of *Lessingianthus*. The statistical analysis showed that presence or absence of idioblasts and crystal types are the most influential variables. These variables, along with the shape and fruit type, allow differentiation among related species. In the genus, there are several complexes of closely related species that are morphologically similar, but with different ploidy levels. The search for additional diagnostic characters to distinguish the species within these groups is necessary, and cypsel characters analysed here contributed to their differentiation. For example, the *L. saltensis* complex is one of these groups that includes three species (*L. saltensis*, *L. coriarius* and *L. membranifolius*), which can be distinguished from each other by morphological features, such as texture and shape of leaves, phyllary apex, chromosome number, as well as by some micromorphological features such as corolla pubescence (Angulo and Dematteis 2012a, 2014). Cypselae were also different in *L. coriarius* where the fruits are glabrous and cylindrical-obconical, whereas *L. saltensis* had obconical fruit with twin hairs and glandular trichomes, and *L. membranifolius* had fruit with only twin hairs. Therefore, the cypsel characters are a useful tool to characterise and distinguish species, and their taxonomic value increases when they are considered together with other morphological characters.

As with some other micromorphological characters of Vernoniae, fruit features of *Lessingianthus* are shared with related genera, namely, *Chrysolaena* and *Lepidaploa*, which hinders their use as good taxonomic markers at a generic level. For example, the absence of glandular trichomes on fruits was previously noted as one of the features that distinguish *Lessingianthus* from *Chrysolaena* and *Lepidaploa* (Robinson 1988a). However, our results indicated that some species of *Lessingianthus* have glandular trichomes; therefore, this character is shared with these related genera. Crystal shape was another characteristic that distinguished *Lessingianthus* (quadrate or subquadrate; designed here as 'cubic or subcubic') from *Lepidaploa* and *Chrysolaena* (elongate, designed here as 'prismatic'; Robinson 1999). However, our results showed a wide variation in crystals, indicating that this character is also not useful for generic segregation. Idioblasts are present in *Lessingianthus* as well as in *Lepidaploa* and *Chrysolaena* (Robinson 1999), so the presence of these cells is shared by the three genera also. Several studies have shown that micromorphological characters are valuable within the genera of Vernoniae and, when combined with other characters (pollen morphology, chromosome number and inflorescence type), they delimit subtribes, genera and species within geographically defined regions (Robinson 1999; Funk *et al.* 2009). When

Robinson (1988a) segregated *Lessingianthus*, he also used a combination of macro- and micromorphological characters. The cypsela features here analysed may not be good taxonomic markers for the genus; however, they are useful at a species-level classification.

Acknowledgements

We appreciate support of this work by grants from the Consejo Nacional de Investigaciones Científicas y Tecnológicas (CONICET), the Secretaría General de Ciencia y Técnica de la Universidad Nacional del Nordeste (SGCyT-UNNE), and the Myndel Botanica Foundation.

References

- Abid R, Alam J (2011) Taxonomic significance of cypsel morphology for the tribe Mutisieae (S.L.) (Asteraceae) from Pakistan. *Pakistan Journal of Botany* **43**, 821–826.
- Abid R, Ali N (2010) Cypsel morphology and its taxonomic significance for the tribe Senecioneae (Asteraceae) from Pakistan. *Pakistan Journal of Botany* **42**, 117–133.
- Abid R, Qaiser M (2009) Taxonomic significance of the cypsel morphology in the tribe Anthemideae (Asteraceae) from Pakistan and Kashmir. *Pakistan Journal of Botany* **41**, 555–579.
- Anderberg AA (1989) Phylogeny and reclassification of the tribe Inuleae (Asteraceae). *Canadian Journal of Botany* **67**, 2277–2296. doi:10.1139/b89-292
- Anderson LE (1954) Hoyer's solution as a rapid mounting medium for bryophytes. *The Bryologist* **57**, 242–247. doi:10.1639/0007-2745(1954)57[242:HSAARP]2.0.CO;2
- Angulo MB, Dematteis M (2010) Pollen morphology of the South American genus *Lessingianthus* (Vernonieae, Asteraceae) and its taxonomic implications. *Grana* **49**, 12–25. doi:10.1080/00173130903435192
- Angulo MB, Dematteis M (2012a) Cytotaxonomy of some species of the South American genus *Lessingianthus* (Asteraceae, Vernonieae). *Plant Systematics and Evolution* **298**, 277–285. doi:10.1007/s00606-011-0542-z
- Angulo MB, Dematteis M (2012b) Taxonomy of the *Lessingianthus saltensis* (Vernonieae, Asteraceae) species complex. *Annales Botanici Fennici* **49**, 239–247. doi:10.5735/085.049.0404
- Angulo MB, Dematteis M (2014) Floral microcharacters in the genus *Lessingianthus* (Vernonieae, Asteraceae) and its taxonomic implications. *Plant Systematics and Evolution* **300**, 1925–1940. doi:10.1007/s00606-014-1019-7
- Bremer K (1994) 'Asteraceae: Cladistic and Classification.' (Timber Press: Portland, OR)
- Bunwong S, Chantaranonthais P, Keeley S (2014) Revisions and key to the Vernonieae (Compositae) of Thailand. *PhytoKeys* **37**, 25–101. doi:10.3897/phytokeys.37.6499
- Cabrera AL (1944) Vernonieae Argentinas (Compositae). *Darwiniana* **6**, 265–379.
- Carpenter KJ (1999) Comparative morphology of disk floret trichomes of *Encelia* (Asteraceae: Heliantheae). PhD thesis, California State Polytechnic University, Pomona.
- Castro MM, Castro H, Leitão-Filho F, Monteiro WR (1997) Utilização de estruturas secretoras na identificação dos gêneros de Asteraceae de uma vegetação do cerrado. *Revista Brasileira de Botânica. Brazilian Journal of Botany* **20**, 163–174. doi:10.1590/S0100-84041997000200007
- Ciccarelli D, Garbari F, Pagni AM (2007) Glandular hair of the ovary: a helpful character of Asteroidae (Asteraceae) taxonomy? *Annales Botanici Fennici* **44**, 1–7.
- Di Rienzo JA, Casanoves F, Balzarini MG, González L, Tablada M, Robledo CW (2013) InfoStat versión 2013. Grupo InfoStat, FCA, Universidad Nacional de Córdoba.
- Dormer KJ (1961) The crystals in the ovaries of certain Compositae. *Annals of Botany* **25**, 241–254.
- Faust WZ, Jones SB Jr (1973) The systematic value of trichome complements in a North American group of *Vernonia* (Compositae). *Rhodora* **75**, 517–528.
- Franceschi VR, Horner HT (1980) Calcium oxalate crystals in plants. *Botanical Review* **46**, 361–427. doi:10.1007/BF02860532
- Freire SE, Katinas L (1995) Morphology and ontogeny of the cypsel hairs of Nassauviinae (Asteraceae, Mutisieae). In 'Advances in Compositae Systematics'. (Eds DJN Hind, C Jeffrey, GV Pope) pp. 107–143. (Royal Botanic Gardens, Kew: London)
- Funk VA, Susanna A, Stuessy T, Robinson H (2009) Classification of Compositae. In 'Systematics, Evolution, and Biogeography of Compositae'. (Eds VA Funk, A Susanna, TF Stuessy, RJ Bayer) pp. 171–189. (International Association of Plant Taxonomy: Vienna, Austria)
- Gonzalez AM, Cristóbal CL (1997) Anatomía y ontogenia de semillas de *Helicteres lhotzkyana* (Sterculiaceae). *Bonplandia* **9**, 287–294.
- Grau L (1977) Astereae: systematic review. In 'The Biology and Chemistry of the Compositae. Vol. 1'. (Eds VH Heywood, JB Harborne, BL Turner) pp. 539–565. (Academic Press: London)
- Haque MZ, Godward MBE (1984) New records of the carpodoid in Compositae and its taxonomic use. *Botanical Journal of the Linnean Society* **89**, 321–340. doi:10.1111/j.1095-8339.1984.tb02564.x
- Hind DJN (1993) Notes on the Compositae of Bahia, Brazil. *Kew Bulletin* **48**, 245–277. doi:10.2307/4117932
- Horner HT (1977) A comparative light and electron microscopic study of microsporogenesis in male-fertile and cytoplasmic male-sterile sunflower (*Helianthus annuus*). *American Journal of Botany* **64**, 745–759. doi:10.2307/2441727
- Ike S, Nordenstam B (2012) Diversity of trichomes from mature cypselar surface of some taxa from the basal tribes of Compositae. *Compositae Newsletter* **50**, 78–124.
- Isawumi MA, El-Ghazaly G, Nordenstam B (1996) Pollen morphology, floral microcharacters and taxonomy of the genus *Baccharoides* Moench (Vernonieae: Asteraceae). *Grana* **35**, 205–230. doi:10.1080/00173139609430008
- Johansen DA (1940) 'Plant Microtechnique.' (McGraw-Hill Book Co.: New York)
- Keeley SC, Forsman ZH, Chan R (2007) A phylogeny of the 'evil tribe' (Vernonieae: Compositae) reveals Old/New World long distance dispersal: support from separate and combined congruent datasets (*trnL-F*, *ndhF*, ITS). *Molecular Phylogenetics and Evolution* **44**, 89–103. doi:10.1016/j.ympev.2006.12.024
- King RM, Robinson H (1970) The new synanthology. *Taxon* **19**, 6–11. doi:10.2307/1217907
- Kraus G (1866) Über die Bau trocken Pericarp. *Jahrbücher für Wissenschaftliche Botanik* **5**, 93–126.
- Lander NS, Hurter PJH (2013) *Pleurocarpaea gracilis* (Asteraceae: Vernonieae), a new species from the Pilbara region of Western Australia. *Nuytsia* **23**, 109–115.
- Luque R, Sousa HC, Kraus JE (1996) Métodos de coloração de Roeser (1972) – modificado – E Kropp (1972), visando a substituição do Azul de Astra por Azul de Alcão 8GS ou 8GX. *Acta Botanica Brasílica* **10**, 199–212.
- Martins MAG, Oliveira DMT (2007) Ontogênese do fruto de *Vernonia brevifolia* e *V. herbacea*. *Revista Brasileira de Botânica. Brazilian Journal of Botany* **30**, 101–112. doi:10.1590/S0100-84042007000100010
- Marzinek J, Cavalari de-Paula O, Trombert Oliveira DM (2008) Cypsel or achene? Refining terminology by considering anatomical and historical factors. *Revista Brasileira de Botânica. Brazilian Journal of Botany* **31**, 549–553. doi:10.1590/S0100-84042008000300018
- Mazen AMA, Zhang D, Franceschi VR (2004) Calcium oxalate formation in *Lemna minor*: physiological and ultrastructural aspects of high capacity

- calcium sequestration. *New Phytologist* **161**, 435–448. doi:[10.1111/j.1469-8137.2004.00923.x](https://doi.org/10.1111/j.1469-8137.2004.00923.x)
- Meric C (2008) Calcium oxalate crystals in *Conyza canadensis* (L.) Cronq. and *Conyza bonariensis* (L.) Cronq. (Asteraceae: Astereae). *Acta Biologica Szegediensis* **52**, 295–299.
- Meric C, Dane F (2004) Calcium oxalate crystals in floral organs of *Helianthus annuus* L. and *H. tuberosus* L. (Asteraceae). *Acta Biologica Szegediensis* **48**, 19–23.
- Metcalfe CR, Chalk L (1979) 'Anatomy of the Dicotyledons. Vol. I.' (Clarendon Press: Oxford, UK)
- Molano-Flores B (2001) Herbivory and calcium concentrations affect calcium oxalate crystal formation in leaves of *Sida* (Malvaceae). *Annals of Botany* **88**, 387–391. doi:[10.1006/anbo.2001.1492](https://doi.org/10.1006/anbo.2001.1492)
- Nakata PA (2003) Advances in our understanding of calcium oxalate crystal formation and function in plants. *Plant Science* **164**, 901–909. doi:[10.1016/S0168-9452\(03\)00120-1](https://doi.org/10.1016/S0168-9452(03)00120-1)
- Narayana BM (1979) Taxonomic value of trichomes in *Vernonia* Schreb. (Asteraceae). *Proceedings of the Indian Academy of Sciences*, **88**(Section A), 347–357.
- Robinson H (1988a) Studies in the *Lepidaploa* complex (Vernonieae: Asteraceae). IV. The new genus *Lessingianthus*. *Proceedings of the Biological Society of Washington* **100**, 929–951.
- Robinson H (1988b) Studies in the *Lepidaploa* complex (Vernonieae: Asteraceae). V. The new genus *Chrysolaena*. *Proceedings of the Biological Society of Washington* **101**, 952–958.
- Robinson H (1999) Generic and subtribal classification of American Vernonieae. *Smithsonian Contributions to Botany* **89**, 1–116. doi:[10.5479/si.0081024X.89](https://doi.org/10.5479/si.0081024X.89)
- Robinson H (2007) Tribe Vernonieae. In 'The Families and Genera of Vascular Plants. Vol. 8'. (Eds J Kadereit, C Jeffrey) pp. 165–192. (Springer: Berlin)
- Robinson H (2009) An introduction to micro-characters of Compositae. In 'Systematics, Evolution and Biogeography of Compositae'. (Eds VA Funk, A Susanna, TF Stuessy, RJ Bayer) pp. 89–100. (International Association for Plant Taxonomy: Vienna, Austria)
- Robinson H, King RM (1977) Eupatorieae-systematic review. In 'The Biology and Chemistry of the Compositae'. (Eds VH Heywood, JB Harborne, BL Turner) vol. 1, pp. 437–485. (Academic Press: New York)
- Roque N, Funk V (2013) Morphological characters add support for some members of the basal grade of Asteraceae. *Botanical Journal of the Linnean Society* **171**, 568–586. doi:[10.1111/boj.12000](https://doi.org/10.1111/boj.12000)
- Roque N, Keil DJ, Susanna A (2009) Illustrated glossary of Compositae. In 'Systematics, Evolution, and Biogeography of Compositae'. (Eds VA Funk, A Susanna, TF Stuessy, RJ Bayer) pp. 781–806. (International Association of Plant Taxonomy: Vienna, Austria)
- Sancho G, Katinas L (2002) Are the trichomes in corollas of Mutisieae (Asteraceae) really twin hairs? *Botanical Journal of the Linnean Society* **140**, 427–433. doi:[10.1046/j.1095-8339.2002.00113.x](https://doi.org/10.1046/j.1095-8339.2002.00113.x)
- Spjut RW (1994) A systematic treatment of fruit types. *Memoirs of the New York Botanical Garden* **70**, 1–182.
- Werker E, Fahn A (1981) Secretory hairs of *Inula viscosa* (L.) Ait. Development ultrastructure and secretion. *Botanical Gazette* **142**, 461–476. doi:[10.1086/337247](https://doi.org/10.1086/337247)
- Werker E, Putievsky E, Ravid U, Dudai N, Katzir I (1994) Glandular hairs, secretory cavities, and the essential oil in leaves of tarragon (*Artemisia dracunculoides* L.). *Journal of Herbs, Spices & Medicinal Plants* **2**, 19–32. doi:[10.1300/J044v02n03_04](https://doi.org/10.1300/J044v02n03_04)
- Zarlavsky GE (2014) 'Histología Vegetal: técnicas simples y complejas.' (Sociedad Argentina de Botánica: Buenos Aires)
- Zhang JW, Boufford DE, Sun H (2013) Systematic significance of achene morphology in *Soro-seris*, *Syncalathium* and *Parasyncalathium* (Asteraceae: Cichorieae). *Botanical Journal of the Linnean Society* **173**, 476–486. doi:[10.1111/boj.12046](https://doi.org/10.1111/boj.12046)

Appendix 1. Specimens analysed

- Lessingianthus adenophyllus* (Mart. ex DC.) H. Rob.
 BRAZIL, Minas Gerais: Menezes *et al.* 11878 (CTES).
L. ammophyllus (Gardner) H. Rob.
 BRAZIL, Minas Gerais: Smith *et al.* 103262 (CTES).
L. arachniolopsis (Ekman & Dusen) H. Rob.
 BRAZIL, Paraná: Barbosa and Costa 1025 (CTES).
L. arctatus Dematt.
 BRAZIL, Goiás: Magenta *et al.* 266 (CTES).
L. argenteus (Less.) H. Rob.
 BRAZIL, Paraná: Hatschbach 1989 (CTES). PARAGUAY, Amambay: Dematteis *et al.* 3396 (CTES).
L. argyrophyllus (Less.) H. Rob.
 BRAZIL, Brasília: Krapovickas *et al.* 31158 (CTES). Minas Gerais: Krapovickas *et al.* 33353 (CTES).
L. asteriflorus (Mart. ex DC.) H. Rob.
 BRAZIL, Paraná: Silva *et al.* 6099 (CTES), BOLIVIA, Santa Cruz: Solís Neffa *et al.* 1351 (CTES).
L. bardanoides (Less.) H. Rob.
 BRAZIL, Mato Grosso do Sul: Hatschbach *et al.* 58736 (CTES). PARAGUAY, Amambay: Dematteis *et al.* 3393 (CTES).
L. barrosoanus Dematt.
 BRAZIL, Bahia: Guédès *et al.* s.n. (SPF).
L. bakerianus Dematt.
 BRAZIL, Minas Gerais: Warming 2867 (CTES).
L. bellulus Dematt.
 BRAZIL, Mato grosso do Sul: Hatschbach *et al.* 58687 (CTES).
L. bishopii (H. Rob.) H. Rob.
 BRAZIL, Goiás: Saint Hilaire 487 (P).
L. brevifolius (Less.) H. Rob.
 ARGENTINA, Corrientes: Angulo 11 (CTES). BRAZIL, Paraná: Silva *et al.* 5152 (CTES). PARAGUAY, Alto Paraná: Schinini *et al.* 28196 (CTES).
 URUGUAY, Rivera: Bonifacino and Wilson 2140 (CTES).
L. brevipetiolatus (Sch. Bip. ex Baker) H. Rob.
 BRAZIL, Minas Gerais: Roth 1615 (CESJ).
L. buddlejifolius (Mart. ex DC.) H. Rob.
 BRAZIL, Goiás: Krapovickas *et al.* 33270 (CTES).
L. carduoides (Baker) H. Rob.
 BRAZIL, Goiás: Gardner 4192 (P).
L. carvalhoi (H. Rob.) H. Rob.
 BRAZIL, Bahia: Conceição 785 (ALCB); Harley *et al.* 50404 (CTES).
L. cataractarum (Hieron.) H. Rob.
 ARGENTINA, Misiones: Gatti 28 (CTES). BRAZIL, Paraná: Ribas *et al.* 6258 (CTES).
L. centauropsideus (Hieron.) Dematt.
 ARGENTINA, Salta: Arbo *et al.* 9107 (CTES).
L. cephalotes (DC) H. Rob.
 BRAZIL, Goiás: Glaziou 21577 (G). Minas Gerais: Mello Silva *et al.* CFCR 5768, SPF 35568 (CTES).
L. clavatus (Gardner) Dematt.
 BRAZIL, Minas Gerais: Mexia 5536 (P).
L. compactiflorus (Mart. ex Baker) H. Rob.
 BRAZIL, Goiás: Hatschbach 43705 (CTES).
L. constrictus (Matzenb. & Mafioletti) Dematt.
 BRAZIL, Rio Grande do Sul: Krapovickas and Cristóbal 44772 (CTES). Irgang 9865 (CTES).
L. cordiger (Mart. ex DC.) H. Rob.
 BRAZIL, Minas Gerais: Hatschbach *et al.* 72053 (MBM, CTES). Semir *et al.* CFCR 9510, SPF 42759 (CTES).
L. coriaceus (Less.) H. Rob.
 BRAZIL, Minas Gerais: Krapovickas *et al.* 33375 (CTES). BOLIVIA, Santa Cruz: Dematteis *et al.* 2066 (CTES).
L. coriarius M.B. Angulo
 BOLIVIA, Chuquisaca: Beck 6252 (CTES). Santa Cruz: Dematteis *et al.* 3660 (CTES); Fuentes 2372 (CTES).
L. correntinus (Cabrera & Cristóbal) Dematt.
 ARGENTINA, Corrientes: Schinini and Ahumada 13895 (CTES).
L. cristalinae (H. Rob.) H. Rob.
 BRAZIL, Goiás: Barroso 522 (UB).
L. durus (Mart. ex DC.) H. Rob.
 BOLIVIA, Santa Cruz: Dematteis *et al.* 2102 (CTES). BRAZIL, Goiás: Hatschbach *et al.* 70856 (CTES).
L. elegans (Gardner) H. Rob.
 BRAZIL, Minas Gerais: Souza *et al.* 10107 B (CTES).
L. exiguus (Cabrera) H. Rob.

(continued next page)

Appendix 1. (continued)

- BRAZIL, Paraná: Krapovickas and Cristobal 40842 (CTES).
L. farinosus (Baker) H. Rob.
 BRAZIL, Bahia: Giulietti *et al.* CFCR 1293 (CTES).
L. floccosus (Gardner) H. Rob.
 BRAZIL, Minas Gerais: Gardner 4786 (G).
L. foliosus Dematt.
 BRAZIL, San Pablo: Glaziou 8133 (P).
L. glabratus (Less.) H. Rob.
 ARGENTINA, Corrientes: Vanni *et al.* 1476 (CTES).
L. grandiflorus (Less.) H. Rob.
 BRAZIL, Goiás: Hatschbach *et al.* 70744 (CTES), Paraná: Krapovickas and Cristobal 40756 (CTES). PARAGUAY, Amambay: Dematteis and Schinini 859 (CTES).
L. grearii (H. Rob.) H. Rob.
 BRAZIL, Goiás: Paula-Souza *et al.* 4563 (CTES).
L. hasslerianus (Chodat) Dematt.
 PARAGUAY, Itapúa: Montes 7161 (CTES).
L. hoveifolius (Gardn.) H. Rob.
 BRAZIL, Goiás: Hatschbach *et al.* 70576 (CTES).
L. hypochaeris (DC.) H. Rob.
 BRAZIL, Paraná: Silva and Ribas 3491 (CTES).
L. hystricosus (Cabrera & Dematt.) Dematt.
 PARAGUAY, Amambay: Dematteis *et al.* 905 (CTES). Cordillera: Schinini 10908 (CTES).
L. ibitipocensis Borges & Dematt.
 BRAZIL, Minas Gerais: Saint Hilaire 200 (P).
L. intermedius (DC.) Dematt.
 URUGUAY, Colonia: Solis Neffa *et al.* 2089 (CTES). Montevideo: Sellow s. n. (BR).
L. irwinii (G.M. Barroso) H. Rob.
 BRAZIL, Goiás: Glaziou s.n. (G).
L. ixiamensis (Rusby) H. Rob.
 BOLIVIA, Beni: Krapovickas and Schinini 34882 (CTES). BRAZIL, Mato Grosso: Pedersen 12213 (CTES).
L. kuntzei (Hieron.) Dematt.
 BOLIVIA, Santa Cruz: Saravia Toledo 12309 (CTES); Vargas 880 (CTES).
L. laevigatus (Mar. ex DC.) H. Rob.
 BOLIVIA, Beni: Solomon 7881 (CTES). BRAZIL, Minas Gerais: Hatschbach *et al.* 64730 (CTES).
L. lanatus (Cabrera) Dematt.
 PARAGUAY, Caaguazú: Schinini *et al.* 36145 (CTES). Cordillera: Schinini 2211 (CTES).
L. laniferus (Cristobal & Dematt.) M. B. Angulo
 ARGENTINA, Corrientes: Krapovickas *et al.* 18056b (CTES). Misiones: Maruñak 119 (CTES). BRAZIL, Santa Catarina: Ribas and Hatschbach 7513 (CTES).
 URUGUAY, Rivera: Dematteis and Schinini 1448 (CTES).
L. lanuginosus Dematt.
 BRAZIL, Brasília: Krapovickas *et al.* 33175 (CTES). Goiás: Serra dos Pireneus, Hatschbach *et al.* 70059 (CTES).
L. lapinhensis Dematt.
 BRAZIL, Minas Gerais: Pirani *et al.* 12129 (CTES).
L. laurifolius (H. B. K.) H. Rob.
 BOLIVIA, La Paz: Beck 19805 (CTES).
L. ligulifolius (Mart. ex DC.) H. Rob.
 BRAZIL, Distrito Federal: Krapovickas and Cristóbal 3190 (CTES). Goiás: Hatschbach and Silva 59996 (CTES).
L. linearifolius (Less.) H. Rob.
 BRAZIL, Minas Gerais: Saavedra *et al.* 270 (CTES). Serra de Ibitipoca: Sucre and Krieger 6850 (CTES).
L. linearis (Spreng.) H. Rob.
 BRAZIL, Bahia: Ganev 186 (CTES). Minas Gerais: Vidal I-828 (CTES).
L. longicuspis Dematt.
 BOLIVIA, Santa Cruz: Dematteis *et al.* 2186 (CTES).
L. lorentzii (Hieron.) H. Rob.
 ARGENTINA, Corrientes: Carnevali 4413 (CTES). Entre Ríos: Martínez Crovetto 4818 (CTES).
L. macrocephalus (Less.) H. Rob.
 BRAZIL, Rio Grande do Sul: Deble *et al.* s.n. (CTES). URUGUAY, Rivera: Pedersen 11648 (CTES).
L. macrophyllus (Less.) H. Rob.
 BRAZIL, Paraná: Hatschbach *et al.* 66520 (CTES).
L. mansoanus (Baker) H. Rob.
 BRAZIL, Mato Grosso do Sul: Hatschbach *et al.* 66643 (CTES). Mato Grosso: Hatschbach *et al.* 66727 (CTES).
L. membranifolius M.B. Angulo

(continued next page)

Appendix 1. (continued)

- BOLIVIA, Santa Cruz: Dematteis *et al.* 2383 (CTES); Dematteis *et al.* 2362 (CTES).
L. minimus Dematt.
 BRAZIL, Goiás: Hatschbach and Kummrow 37253 (CTES).
L. mollissimus (D. Don ex Hook & Arn.) H.Rob.
 ARGENTINA, Misiones: Krapovickas and Cristóbal 28841 (CTES). BRAZIL, Rio Grande do Sul: Hagelund 7851 (CTES). PARAGUAY, Alto Paraná: Caballero Marmori s.n. (CTES).
L. monocephalus (Gardner) H. Rob.
 BRAZIL, Goiás: Pereira-Silva *et al.* 9935 (CTES), Minas Gerais: Schinini and Barbosa 71330 (CTES).
L. morii (H.Rob.) H.Rob.
 BRAZIL, Bahia: Hatschbach *et al.* 67666 (CTES).
L. myrsinites H.Rob.
 BRAZIL, Goiás: Hatschbach *et al.* 60128 (CTES).
L. niederleinii (Hieron.) H.Rob.
 ARGENTINA, Misiones: Montes 192 (CTES). BRAZIL, Mato Grosso do Sul: Hatschbach *et al.* 74473 (CTES). PARAGUAY, Caaguazú: Lopez *et al.* 196 (CTES).
L. obscurus (Less.) H.Rob.
 BRAZIL, Goiás: Gardner 3791 (G).
L. obtusatus (Less.) H.Rob.
 BOLIVIA, Santa Cruz: Killeen *et al.* 6499 (CTES). BRAZIL, Goiás: Hatschbach *et al.* 60116 (CTES).
L. onopordioides (Baker) H.Rob.
 BRAZIL, Mato Grosso do Sul: Souza *et al.* 16972 (CTES).
L. parvifolius (Chodat) H.Rob.
 PARAGUAY, Cordillera: Bordas 1257 (CTES). Paraguari: Mereles *et al.* 8453 (CTES).
L. pentacontus (DC) H.Rob.
 BRAZIL, Minas Gerais: Hatschbach *et al.* 64852 (CTES); Hatschbach *et al.* 64644 (CTES).
L. plantaginodes (Kuntze) H.Rob.
 ARGENTINA, Corrientes: San Martín. Medina and Salas 378 (CTES). Entre Ríos: Krapovickas and Cristobal 46566 (CTES). BRAZIL, Porto Alegre: Bueno 2148 (CTES). URUGUAY, Soriano: Castellanos 17612 (CTES).
L. platyphyllus (Chodat) H.Rob.
 BRAZIL, Mato Grosso do Sul: Hatschbach *et al.* 74558 (CTES). PARAGUAY, Amambay: Dematteis and Schinini 865 (CTES).
L. polyphyllus (Sch. Bip. ex Baker) H.Rob.
 ARGENTINA, Misiones: Dematteis 2752 (CTES). BRAZIL, Paraná: Hatschbach 79600 (CTES). PARAGUAY, Amambay: Schinini *et al.* 36102 (CTES).
L. profusus (Dematt. & Cabrera) M. B. Angulo
 BRAZIL, San Pablo: Panizza SPF 17789 (CTES). PARAGUAY, Amambay: Schinini *et al.* 30440 (CTES).
L. pseudoincanus (Hieron.) Dematt. & Angulo
 ARGENTINA, Buenos Aires: Mazzucconi 967 (CTES). Corrientes: Angulo 9 (CTES). Salta: Tolaba 2039 (CTES). BRAZIL, Rio Grande do Sul: Rambo 1968 (LP). URUGUAY, Rocha: Rossengurt 2448 (LP).
L. pseudopiptocarphus (H.Rob.) H.Rob.
 BRAZIL, Goiás: Machado Teles *et al.* 1906 (RB).
L. psilophyllus (Sch. Bip. ex Baker) H.Rob.
 BRAZIL, Minas Gerais: Hatschbach *et al.* 51076 (CTES); Hatschbach *et al.* 64716 (CTES).
L. pubescens Angulo & Dematt.
 BOLIVIA, Boquerón: Vanni *et al.* 2418 (CTES). PARAGUAY, Amambay: Dematteis and Schinni 876 (CTES, G).
L. pumilus (Vell.) H.Rob.
 BRAZIL, Paraná: Hatschbach 25932 (CTES)
L. pusillus (Dematt.) M.B. Angulo.
 ARGENTINA, Chaco: Dematteis 605 (CTES). Corrientes: Dematteis *et al.* 2769 (CTES). Formosa: Morel 5107 (CTES). PARAGUAY, Concepción: Dematteis *et al.* 3234 (CTES).
L. pycnostachys (DC) H.Rob.
 BRAZIL, Minas Gerais: Motta 1243 (CTES); Arbo *et al.* 4949 (CTES).
L. ramellae (Cabrera) H.Rob.
 PARAGUAY, Paraguari: Hassler 6674 (G).
L. regis (H.Rob.) H.Rob.
 BRAZIL, Minas Gerais: Zappi *et al.* 10301 (CTES).
L. reitzianus (Cabrera) H.Rob.
 BRAZIL, Santa Catarina: Silva and Andrade 5462 (CTES). Paraná: Ferrucci *et al.* 236 (CTES); Hatschbach 39206 (CTES).
L. robustus (Rusby) H.Rob.
 BOLIVIA, La Paz: Krapovickas 46754 (CTES). Santa Cruz: Fuentes and Navarro 2183 (CTES). BRAZIL, Mato Grosso: Riedel 1474 (P).
L. roseus (Mart. ex DC.) H.Rob.
 BRAZIL, Minas Gerais: Krapovickas and Cristobal 33517 (CTES); Abudd *et al.* CFSC 10752 (CTES); Damazio 1465 (RB).
L. rosmarinifolius (Less.) H.Rob.

(continued next page)

Appendix 1. (continued)

-
- BRAZIL, Minas Gerais: Arbo *et al.* 3913 (CTES); Arbo *et al.* 4170 (CTES).
L. rubricaulis (Humb. & Bonpl.) H.Rob.
 ARGENTINA, Corrientes: Dematteis *et al.* 2756 (CTES). Misiones: Dematteis and Surenciski 2452 (CTES). BOLIVIA, Santa Cruz: Dematteis *et al.* 3567 (CTES). BRAZIL, Mato Grosso do Sul: Hatschbach *et al.* 74696 (CTES). PARAGUAY, Amambay: Dematteis *et al.* 3375 (CTES).
L. saltensis (Hieron.) H.Rob.
 ARGENTINA, Jujuy: Dematteis *et al.* 2952 (CTES). Salta: Pozner and Belgrano 454 (CTES). BOLIVIA, Chiquitos: Dematteis *et al.* 2334 (CTES, SI).
L. sancti-pauli (Hieron.) Dematt.
 BRAZIL, Paraná: Von Lisingen and Sonehara 115 (CTES).
L. santosii (H.Rob.) H.Rob.
 BRAZIL, Bahia: Giulietti 1275 (CTES). Minas Gerais: Arbo *et al.* 5133 (CTES).
L. scabrifolius (Hieron.) H.Rob.
 BOLIVIA, Santa Cruz: Seijo 3215 (CTES).
L. secundus (Sch. Bip. ex Baker) H.Rob.
 BRAZIL, Goiás: Glaziou 21634 (R).
L. sellowii (Less.) H.Rob.
 ARGENTINA, Misiones: Dematteis 588 (CTES). BRAZIL, Rio Grande do Sul: Deble and Oliveira, *s.n.* (CTES). URUGUAY, Tacuarembó: Dematteis *et al.* 3760 (CTES).
L. soderstroemii (H.Rob.) H.Rob.
 BRAZIL, Minas Gerais: Souza *et al.* 25168 (ESA); Hatschbach *et al.* 54521 (CTES); Hatschbach *et al.* 58357 (CTES).
L. souzae (H.Rob.) H.Rob.
 BRAZIL, Goiás: Glaziou 21632 (BR).
L. spicatus (Cabrera) Dematt.
 ARGENTINA, Misiones: Schwatz 5491 (CTES); Schulz 185 (CTES).
L. stoechas (Mart. ex DC.) H.Rob.
 BRAZIL, Minas Gerais: Hatschbach *et al.* 72051 (CTES).
L. subobscurus (Malme) H.Rob.
 BOLIVIA, Santa Cruz: Seijo and Solis Neffa 3248 (CTES).
L. syncephalus (Sch. Bip.) H.Rob.
 BRAZIL, San Pablo: Hohene 2102 (CTES). PARAGUAY, Alto Paraná: Schinini *et al.* 28185 (CTES).
L. teyucuaensis (Cabrera) Dematt.
 ARGENTINA, Misiones: Dematteis *et al.* 515 (CTES).
L. tomentellus (Mart. ex DC.) H.Rob.
 BRAZIL, Minas Gerais: Hatschbach *et al.* 64579 (CTES).
L. ulei (Hieron.) H.Rob.
 BRAZIL, Paraná: Barbosa and Cordeiro 1195 (CTES); Ribas *et al.* 5060 (CTES).
L. varroniiifolius (DC.) H.Rob.
 BOLIVIA, Santa Cruz: Dematteis *et al.* 3860 (CTES); Beck 25852 (CTES).
L. venosissimus (Sch. Bip. Ex Baker)
 BRAZIL, Mato Grosso: Hatschbach *et al.* 66618 (CTES). Rondonia: Krapovickas *et al.* 40142 (CTES).
L. vepretorum (Mart. ex DC.) H.Rob.
 BRAZIL, Minas Gerais: Weddell 1073 (G); Pirani CFSC 12815 (CTES).
L. vestitus (Baker) H.Rob.
 BRAZIL, Minas Gerais: Claussen s. n. (P).
L. warmingianus (Baker) H.Rob.
 BRAZIL, Minas Gerais: Glaziou 20383 (R); Krapovickas and Cristobal 33496 (CTES); Souza *et al.* 29576 (CTES).
L. westermanii (Ekman & Dusén) H.Rob.
 BRAZIL, Paraná: Dusén 16400 (G).
L. xanthophyllus (Mart. ex DC.) H.Rob.
 BRAZIL, Tocantins: Paula-Souza *et al.* 4763 (CTES).
L. zuccarinianus (Mart. ex DC.) H.Rob.
 BRAZIL, Goiás: Weddell 2060 (P). Mato Grosso: Hatschbach *et al.* 62350 (CTES).
-