



## A new polyploid species of *Mecardonia* (Gratioleae, Plantaginaceae) from South America

JULIÁN A. GREPPI<sup>1</sup>, MARÍA DE LAS MERCEDES SOSA<sup>2,3\*</sup> & MASSIMILIANO DEMATTEIS<sup>2,3</sup>

<sup>1</sup>Instituto de Floricultura, Instituto Nacional de Tecnología Agropecuaria, De los Reseros y Las Cabañas s/n°, Hurlingham 1686, Buenos Aires, Argentina.

<sup>2</sup>Instituto de Botánica del Nordeste (UNNE-CONICET), Casilla de Correo 209, W3400CBL Corrientes, Argentina.

<sup>3</sup>Facultad de Ciencias Exactas y Naturales y Agrimensura, Universidad Nacional del Nordeste, Av. Libertad 5460, 3400, Corrientes, Argentina.

\*Author for correspondence. E-mail: [mdlmsvg@yahoo.com.ar](mailto:mdlmsvg@yahoo.com.ar)

### Abstract

*Mecardonia reneae*, a new species from northeastern Argentina, is described and illustrated. This taxon has certain resemblance to *M. procumbens* var. *flagellaris*, but can be easily distinguished by the combination of life form, aspect of the plant, root and stem type, sepal width, fruit size, and chromosome number ( $2n=4x=44$ ). A key to distinguish it from related species is also provided.

### Introduction

*Mecardonia* Ruiz & Pavon (1794: 95) is a New World genus of the tribe Gratioleae (Plantaginaceae). It is characterized by annual or perennial herbs with erect or creeping habit, densely branched, mostly glabrous, glandular-punctate leaves, pedicels basally bibracteolate, yellow and white flowers, and anther thecae separated by an arm-like connective tissue (D'Arcy 1979). In the review by Rossow (1987), 10 species of *Mecardonia* distributed from the east coast of the United States to Argentinian Patagonia and Central Chile were considered. This number of species was modified by Souza (1997) and Souza & Giulietti (2009), who suggested seven species and established the center of diversification including Rio Grande do Sul (Brazil), Uruguay and northeastern Argentina. Recently, another species from Argentina was described: *M. kamogawae* Greppi & Hagiwara (2011: 44). However further studies are needed to clearly delimit the *M. procumbens* complex.

The chromosome number of *Mecardonia* species revealed that the genus has diploid and polyploid taxa (Sosa *et al.* 2016). *Mecardonia grandiflora* (Bentham 1836: 56) Pennell (1946: 87), *M. procumbens* (Miller 1768: 6) Small (1903: 1338) var. *procumbens* and *M. procumbens* var. *flagellaris* (Chamisso & Schlechtendal 1827: 575) Souza (1997: 186) are diploid ( $2n=22$ ), whereas *M. kamogawae* is hexaploid ( $2n=66$ ). Polyploidization has been recognized as an important process in plant evolution (Stebbins 1971, Grant 1989, Otto & Whitton 2000, Soltis *et al.* 2004). Polyploidy is a common phenomenon present in genera of Plantaginaceae, such as *Antirrhinum* Linnaeus (1753: 612), *Cymbalaria* Hill (1756: 113), *Chelone* Linnaeus (1753: 611), *Digitalis* Linnaeus (1753: 621), *Linaria* Miller (1754: 2), *Plantago* Linnaeus (1753: 112), *Stemodia* Linnaeus (1759: 1118), and *Veronica* Linnaeus (1753: 9) (Hair 1966, Subramanian & Pondmudi 1987, Sosa & Seijo 2002, Sosa *et al.* 2009, 2011, Wolfe *et al.* 2002, Vargas *et al.* 2004, Murray *et al.* 2010, Castro *et al.* 2012, Wong & Murray 2012, Ranjbar & Nouri 2015). Previous studies show that in *Stemodia*, diploids and polyploids are quite different in morphological and anatomical features (Sosa *et al.* 2012, Sosa & Dematteis 2014).

The results of a comparative morphological and cytological analysis in *Mecardonia*, allowed us to identify *Mecardonia reneae* as a new species, which is here described. Additionally, an identification key including morphologically similar species is provided.

## Materials and methods

### Field study and plant collection

Material of the new species was collected in the field; part of this material was prepared for herbarium collection, whereas some fruit and flower samples were stored for cytological and micromorphological studies. In addition, some plants were grown in greenhouses. Herbarium material is deposited at BAB, SI.

### Cytological studies

Mitotic chromosome preparations were made from root meristems. The roots were pretreated in 0.002 M 8—hydroxyquinoline solution at room temperature for about 3 h, fixed in 5:1 absolute alcohol/lactic acid (Fernández 1973) and kept in 70% ethanol at 4° C until analysis. Roots tips were stained following Feulgen's technique, and meristems were macerated and squashed in a drop of lactopropionic orcein (Dyer 1963). Permanent microscope slides were prepared in Euparal using the method of Bowen (1956) and examined and photographed under a Carl Zeiss microscope equipped with a Canon Power Shot A 640 digital camera.

## Taxonomic treatment

*Mecardonia reneae* Greppi & Sosa, *sp. nov.* (Figs. 1–2, Table 1)

Type:—ARGENTINA. Corrientes. Dep. Empedrado, Ruta Provincial 6 y arroyo Empedrado [27° 48'S, 58°30'W], 20 December 2005.

Greppi & Hagiwara 560 (holotype BAB!, isotype SI!).


*Diagnosis*:—*A Mecardonia procumbens* var. *flagellaris* similis, sed habitu terophyto, caulibus erectis, radicibus gemmiferis, foliis serratis, petalis angustis, deltoideis et linearibus, fructus minore, chromosomatum numero tetraploideo  $2n=44$  (contra  $2n=22$ ).

Terophyte, 15–17(–20) cm high. Roots gemiferous. Stems erect, quadrangular, glabrous, 1 mm diam., internodes 5–20 mm long. Leaves opposite, sessile, narrowly elliptic, 6–16(–20) × 2–5(–6) mm, margin serrate, base attenuate. Bracteoles 3–7 × 0.5–1.5 mm, linear, margin entire to denticulate. Pedicels 10–35(–40) mm long, erect, flexuous. Flowers one or two per node, 10–16 mm long, yellow with chestnut streaks in the throat. Calyx 5–7 mm long; sepals narrowly deltoid to linear, acute, dorsal sepal 5–7(–10) × 1–1.2 mm; two ventral sepals 5.5–7(–8) × 0.8–1(–3) mm; two lateral sepals 4–6(–8) × 0.3–0.6 mm. Corolla 8–14 × 6–8 mm, posterior lip 6–8 × 6–7 mm, entire, briefly apiculate, internally pubescent at base; middle lobe of frontal lip 6–7 × 7–8.5 mm; broadly obovate, glabrous; lateral lobes of equal size, sometimes obovate. Stamens 4, anterior filaments 5 mm long, posterior 4 mm long, all inserted at the base and reaching half of the corolla tube length. Ovary ellipsoid, 3 mm long; style 3–4 mm long. Capsule ellipsoid, light brown, 3–5 × 1.5–1.8 mm. Seeds, 0.4–0.5 × 0.3–0.4 mm, rough, blackish.


**Distribution, habitat and phenology**:—*Mecardonia reneae* grows in open and humid areas, low clayey loam soils near watercourses, or border of streams in eastern Formosa, Chaco, northwest of Corrientes and northeast of Santa Fe in Argentina, within the Chaco phytogeographic province (Cabrera 1976). The species flowers from August to March.

**Etymology**:—The specific epithet is dedicated to Renée H. Fortunato, Argentinean botanist dedicated to the study of Leguminosae.

**Additional specimens examined**:—ARGENTINA. Prov. Chaco: Dep. Bermejo, Las Palmas, 14 November 1983, Fortunato *et al.* 607 (BAB); Dep. Primero de Mayo, Colonia Benítez, 10 June 1959, Schulz 10437 (BAB). Prov. Corrientes: Dep. Capital, near to Ingenio Primer Correntino, 18 Km NE of Corrientes, 15 September 1982, Schinini & Martínez-Crovetto 2636 (CTES); 5 Km E of Laguna Brava, 29 August 1970, Krapovickas & Cristóbal 15836 (CTES); 10 Km E of Laguna Brava, route 5, 21 August 1970, Krapovickas & Cristóbal 16099 (CTES); Parques de la Facultad de Ciencias Agrarias, 23 August 1974, Quarín 2403 (CTES); Molina Punta, November 1978, Arbo 1562 (CTES); Perichón, 8 September 1975, Schinini 11601 (CTES); Dep. San Cosme, Paso de la Patria, 12 August 1967, Krapovickas & Cristóbal 1302 (CTES); Desvío a Pto. González, 9 Km. near to Paso de la Patria, 10 September 1971, Tressens *et al.* 168 (CTES); Dep. San Luis del Palmar, 10 Km SE of San Luis del Palmar, Route 6, 26 September 1973, Quarín & Tressens 1349 (CTES). Prov. Formosa: Dep. Pilcomayo, Parque Nacional Pilcomayo, near to Puerto Soledad, 9 November 1991, Fortunato *et al.* 2097 (BAB, SI). Prov. Santa Fe: Dep. Gral. Obligado, Villa Ana, 18 August 1973, Quarín 1193 (CTES).



**FIGURE 1.** *Mecardonia reneae*. A: Habit. B: Stem. C: Detail of the leaf. D: Flower in lateral view. E: Calyx. F: Dissected corolla. G: Anterior anther. H: Posterior anther. I: Gynoecium. J: Fruit (A-C: from Greppi & Hagiwara 560, BAB; illustrated by M. Marino; B-J: from Schinini 11601, CTES; illustrated by Mirtha L. Gómez).



**FIGURE 2.** A, C: *Mecardonia procumbens* var. *flagellaris*. A. Habit, C. Metaphase image showing 22 chromosomes. B, D. *M. reneae*. B. Habit. D. Metaphase image showing 44 chromosomes.

**TABLE 1.** Comparison of diagnostic characters between *Mecardonia procumbens* var. *flagellaris*, *M. reneae* and *M. kamogawae*.

	<i>M. procumbens</i> var. <i>flagellaris</i>	<i>M. reneae</i>	<i>M. kamogawae</i>
Life forms	camephyte decumbent	terophyte	geophyte
Stems	repens, radicant, stoloniferous	not radicant	not radicant
Root	not gemmiferous	gemmiferous	gemmiferous
<b>Leaf margin</b>	entire	serrate	serrate
Form leaf	linear to narrow elliptical	narrow elliptic	ovate to broadly elliptic
Consistency of the leaf	herbaceous	herbaceous	slightly succulent
Bracteoles (mm)	5–11 × 0.5–2	3–7 × 0.5–1.5	6–12 × 2–4
Dorsal sepal width (mm)	1.5–3	1–1.2	1.5–2
Capsule size (mm)	5–9 × 2–4	3–5 × 1.5–1.8	5–6 × 3–3.5
Chromosome numbers	2n=22	2n=44	2n=66

## Discussion

We confirmed a chromosome number of  $2n=4x=44$  in *Mecardonia reneae* (Fig. 2), as reported with the name *Mecardonia* sp. in Sosa *et al.* (2016). This taxon differs from other similar taxa that are diploid ( $2n=2x=22$ ), such as *M. procumbens* var. *flagellaris*, or hexaploid, such as *M. kamogawae* ( $2n=2x=66$ ).

Morphologically, *Mecardonia reneae* is similar to *M. kamogawae* in terms of the gemiferous root system (Fig. 2) but can be distinguished by the sepal width, consistence and color of leaves, bracteole length and fruit size. Furthermore, some materials of this new species were mistaken for *M. procumbens* var. *flagellaris* in the past. However, both species can be easily distinguished when studying living plants. *Mecardonia procumbens* var. *flagellaris* are plants with radicating stems, occasionally somewhat ascending at the ends; the stems and its branches grow up to 40 or 50 cm in length, spreading across the ground and producing adventitious roots; it does not have gemiferous roots; its appearance is not compact. In addition, the combination of erect and non-radicant stems and gemmiferous roots gives *M. reneae* a compact appearance. The margin leaves of *M. procumbens* var. *flagellaris* are entire or sub-serrate and the sepals, mainly the dorsal sepal, are wider (1.5–3 mm vs. 1–1.2 mm in width), and the fruits are bigger ( $5-9 \times 2-4$  mm vs.  $3-5 \times 1.5-1.8$  mm) than those of *M. reneae*.

These species were also found to differ in their life form. During the winter season, when visiting different natural populations of *Mecardonia*, we observed that *M. procumbens* var. *flagellaris* maintains aerial shoots latent, whereas *M. reneae* plants disappear completely and grows again when the new spring season approaches. According to the classification of Raunkiaer (1934, modified by Ellenberg & Mueller-Dombois, 1966), *M. procumbens* var. *flagellaris* would behave as a camephyte species, whereas *M. reneae* would be terophyte.

The results obtained from the intra- and inter-specific crosses in native *Mecardonia* species indicated that in the intraspecific crosses *M. reneae* produced 100% of fruits with a high percentage of viable seeds. However, the interespecific crosses with *M. procumbens* var. *procumbens*, *M. procumbens* var. *flagellaris*, *M. kamogawae* and *M. grandiflora* produced no capsules. According to this result, *M. reneae* is reproductively isolated from *M. procumbens* var. *procumbens*, *M. procumbens* var. *flagellaris*, *M. grandiflora* and *M. kamogawae*, and is therefore considered a valid species. The information of crosses is being processed for immediate publication.

There is also a distinction in the distribution among these three species. *Mecardonia reneae* is restricted to northern Argentina; it grows in Chaco province (Cabrera & Willink 1973, Cabrera 1976), in low clayey loam soils near watercourses in areas near Paraná River. *Mecardonia kamogawae* is endemic to the center-east of Corrientes province; it grows in sandy soils, near Uruguay River. In contrast, *M. procumbens* var. *flagellaris* is widely distributed from Mato Grosso do Sul (Brazil) to Chubut (Argentina), including areas in Chile, Paraguay and Uruguay, and covering more phytogeographic provinces. This pattern of cytotype distribution is characteristic of young polyploid complexes (Stebbins 1971). In these complexes, the geographic ranges of polyploid cytotypes are more restricted than those of diploids, but their ranges overlaps with those of the diploid species; thus, this overlap might contribute genomes to the tetraploid species, determining some degree of sympatry among these species.

In conclusion, *M. reneae* seems to be closely related to *M. kamogawae* and in the past some materials of this new species were mistaken for *M. procumbens* var. *flagellaris*. We found that the combination of the morphological and chromosome characters, with geographical range information provides a satisfactory identification of nearly all specimens.

### Key to *M. reneae* and similar species

1. Stems prostrate, radicants, stoloniferous, sometimes with erect or ascending branches born from the prostrate stems. Root system not gemmiferous. Leaves entire. .... *M. procumbens* var. *flagellaris*  
- Stems erect, not radicants, gemmiferous root system. Leaves serrate. .... 2
2. Leaves herbaceous, narrowly elliptical, concolorous. Bracteoles  $3-5 \times 0.5-1.5$  mm. Pedicels up to 35(–40) mm. Dorsal sepal 1–1.2 mm wide. Capsule  $3-5 \times 1.5-1.8$  mm. .... *M. reneae*  
- Leaves slightly succulent, ovate to broadly elliptical, discolorous. Bracteoles  $6-12 \times 2-4$  mm. Pedicels up to 65 mm. Dorsal sepal 1.5–2 mm wide. Capsule  $5-6 \times 3-3.5$  mm ..... *M. kamogawae*

## Acknowledgements

The first author thanks the Direction of Natural Resources of the Corrientes province, also to Tomohiro Kamogawa and Juan Carlos Hagiwara for the support during the Collection trip when the type specimen was collected. We would especially like to thank the keepers and staff of the visited herbaria for their collaboration. The drawing of the species was prepared by Mirtha Liliana Gómez of the Instituto de Botánica del Nordeste and María Angélica Marino of the Instituto Nacional de Tecnología Agropecuaria. This work has been supported by grants from the Consejo Nacional de Investigaciones Científicas y Tecnológicas (CONICET), and the Secretaría General de Ciencia y Técnica of the Universidad Nacional del Nordeste (SGCyT-UNNE).

## References

- Bentham, G. (1836). Observations on some new, or little known genera and species of Scrophulariaceae. *Companion to the Botanical Magazine* 2: 56.
- Bowen, C.C. (1956) Freezing by liquid carbon dioxide in making slides permanent. *Stain Technology* 31: 87–90.  
<http://dx.doi.org/10.3109/10520295609113782>
- Cabrera, A.L. (1976) *Regiones fitogeográficas argentinas*. Ed. Acme, Buenos Aires, 85 pp.
- Cabrera, A.L. & Willink, A. (1973) *Biogeografía de América Latina*.—Monografía N° 13. Secretaría General de la Organización de los Estados Americanos, Programa Regional de Desarrollo Científico y Tecnológico, Washington, DC (USA), 120 pp.
- Castro, M., Castro, S. & Loureiro, J. (2012) Genome size variation and incidence of polyploidy in Scrophulariaceae sensu lato from the Iberian Peninsula. *AoB PLANTS* 2012: pls037.  
<https://doi.org/10.1093/aobpla/pls037>
- Chamisso, A. & Schlechtendal, D. (1827) De plantis in expeditione speculatoria Romanzoffiana observatis disserere pergunt. Scrophularineae. *Linnaea* 2: 555–609.
- Ellenberg, H. & Mueller-Dombois, D. (1966) A key to Raunkiaer plant life forms with revised subdivisions. *Berichte des Geobotanischen Institutes der Eidgenössischen Hochschule, Stiftung Rübli* 37: 56–73.
- D'Arcy, W.G. (1979) Scrophulariaceae. In: Woodson, R.R. & Schery, R.W. (Eds.) *Flora de Panamá*. Parte IX. *Annales of Missouri Botanical Garden* 66: 173–274.
- Dyer, A.F. (1963) The use of lactopropionic orcein in rapid squash methods for chromosome preparations. *Stain Technology* 38: 85–90.
- Fernández, A. (1973) El ácido láctico como fijador cromosómico. *Boletín de la Sociedad Argentina de Botánica* 15: 287–290.
- Grant, V. (1989) *Especiación vegetal*. Limusa, Mexico, D.F., 587 pp.
- Greppi, J.A. & Hagiwara, J.C. (2011) Una nueva especie de *Mecardonia* (Plantaginaceae). *Darwiniana* 49: 43–46.
- Hair, J.B. (1966) Biosystematics of the New Zealand flora, 1945–1964. *New Zealand Journal of Botany* 4: 559–595.
- Hill, J. (1756) *The british herbal*. T. Osborne and J. Shipton, London, 532 pp. Available from: <http://biodiversitylibrary.org/page/34898152> (accessed 1 April 2017)
- Raunkier, C. (1934) *The Life Forms of Plants and Statistical Plant Geography*. Clarendon Press, Oxford, 632 pp.
- Linnaeus C (1753) *Species plantarum*. Salvius, Stockholm, 560 pp.
- Linnaeus C (1759) *Systema Naturae*, Salvius, Stockholm, 984 pp.
- Miller, P (1754) *The Gardener's Dictionary Abridged*. Fourth Edition. P. Miller, London, without pagination.
- Miller, P (1768) *The Gardener's Dictionary*. Eighth Edition. P. Miller, London, without pagination.
- Murray, B.G., Meudt, H.M., Tay, M.L. & Garnock-Jones, P.J. (2010) New chromosome counts in New Zealand species of *Plantago* (Plantaginaceae). *New Zealand Journal of Botany* 48: 197–204.  
<https://doi.org/10.1080/0028825x.2010.515598>
- Otto, S.P. & Whitton, J. (2000) Polyploid incidence and evolution. *Annual Review of Genetics* 34: 401–437.  
<https://doi.org/10.1146/annurev.genet.34.1.401>
- Pennell, F.W. (1946) Reconsideration on the *Bacopa-Herpestis* problem of the Scrophulariaceae. *Proceedings of the Academy of Natural Sciences of Philadelphia* 98: 83–98.
- Ranjbar, M. & Nouri, S. (2015) Biogeography of the genus *Linaria* (Plantaginaceae) based on chromosome number data. *Journal of Cell and Molecular Research* 7: 115–132.  
<https://doi.org/10.22067/jcmr.v7i2.41375>
- Rossow, R.A. (1987) Revisión del género *Mecardonia* (Scrophulariaceae). *Candollea* 42: 431–474.
- Ruiz, L.H. & Pavon, J.A. (1794) *Prodromus flora peruviana et Chilensis*. Sancha, Spain, 116 pp.

- Small, J.K. (1903) *Flora of the Southeastern United States*. J.K. Small, New York, 1370 pp.
- Soltis, D.E., Soltis, P.S. & Tate, J.A. (2004) Advances in the study of polyploidy since plant speciation. *New Phytologist* 161: 173–191.  
<https://doi.org/10.1046/j.1469-8137.2003.00948.x>
- Sosa, M.M. & Seijo, G. (2002) Chromosome Studies in Argentinean Species of *Stemodia* L. (Scrophulariaceae). *Cytologia* 67: 261–266.  
<http://doi.org/10.1508/cytologia.67.261>
- Sosa, M.M., Seijo, G. & Fernández, A. (2009) Chromosome studies in South-American species of *Stemodia* (Scrophulariaceae) and their geographical distribution. *Annales Botanici Fennici* 46: 389–396.  
<http://dx.doi.org/10.5735/085.046.0503>
- Sosa, M.M., Panseri A.F. & Fernández, A. (2011) Karyotype analysis of the southernmost South American species of *Stemodia* (Scrophulariaceae). *Plant Biosystems* 145: 472–477.  
<http://dx.doi.org/10.1080/11263504.2011.566250>
- Sosa, M.M., Panseri, A.F. & Dematteis, M. (2012) Morphometric analysis of *Stemodia hyptoides* and *S. stricta* (Plantaginaceae). *Plant Systematic & Evolution* 298: 1315–1323.  
<https://doi.org/10.1007/s00606-012-0638-0>
- Sosa, M.M. & Dematteis, M. (2014) *Stemodia diplohyptoides* (Plantaginaceae, Gratiolae) a new diploid species from the South America. *Phytotaxa* 186: 271–278.  
<http://dx.doi.org/10.11646/phytotaxa.186.5.4>
- Sosa, M.M., Angulo M.B., Greppi, J.A. & Bugallo, V. (2016) Chromosome numbers and DNA content in some species of *Mecardonia* (Gratiolae, Plantaginaceae). *Comparative Cytogenetic* 10: 769–780.  
<https://doi.org/10.3897/CompCytogen.v10i4.10362>
- Souza, V.C. (1997) Considerações sobre a delimitação de *Mecardonia procumbens* (Mill.) Small (Scrophulariaceae). *Acta Botanica Brasilica* 11: 181–189.  
<http://dx.doi.org/10.1590/S0102-33061997000200008>
- Souza, V. & Giulietti, A.M. (2009) Levantamento das espécies de Scrophulariaceae *sensu lato* nativas do Brasil. *Pesquisas Botânica* 60: 7–288.
- Stebbins, G.L. (1971) *Chromosomal evolution in higher plants*. Arnold, London, 216 pp.
- Subramanian, D. & Pondmudi, R. (1987) Cytotaxonomical studies of South Indian *Scrophulariaceae*. *Cytologia* 52: 529–541.  
<https://doi.org/10.1508/cytologia.52.529>
- Vargas, P.J., Rosselló, J.A., Oyama, R. & Güemes, J. (2004) Molecular evidence for naturalness of genera in the tribe Antirrhineae (Scrophulariaceae) and three independent evolutionary lineages from the New World and the Old. *Plant Systematics and Evolution* 249: 151–172.  
<https://doi.org/10.1007/s00606-004-0216-1>
- Wolfe, A.D., Datwyler, S.L. & Randle, C.P. (2002) A phylogenetic and biogeographic analysis of the Cheloneae (Scrophulariaceae) based on ITS and matK sequence data. *Systematic Botany* 27: 138–148.  
<http://dx.doi.org/10.1043/0363-6445-27.1.138>
- Wong, C. & Murray, B.G. (2012) Variable changes in genome size associated with different polyploid events in *Plantago* (Plantaginaceae). *Journal of Heredity* 103: 711–719.  
<https://doi.org/10.1093/jhered/ess049>