

Thermal Decomposition of Cyclic GlutaraldehydeDiperoxide.UV.Spectroscopic Analysis

A.G. Bordón¹, M. J. Jorge¹, M. I. Profeta¹, J.M. Romero¹, N. L. Jorge¹ and E. A. Castro²

¹LabInTam, Facultad de Cs. Exactas y Naturales y Agrimensura – Universidad Nacional del Nordeste-Av. Libertad 5470 - (3400) Corrientes-Argentina

²INIFTA, TheoreticalChemistryDivision, Suc. 4, C.C. 16, La Plata 1900, Buenos Aires, Argentina

Corresponding Author E-mail: nelly.jorge@gmail.com

Accepted: April 08, 2015

Abstract: The thermal decomposition reaction of Cyclic GlutaraldehydeDiperoxide (3,6-dibutanal-1,2,4,5-tetroxane) in solution of methanol in the temperature range of 130.0-166.0°C and initial concentration of 1.69×10^{-3} mol/L, follows first-order kinetic laws. Under the experimental conditions, the activation parameters of the reaction are bigger than the obtained ones for other tetroxanos with different substituent's in the same solvent, which indicates us that the substituents stabilize the condition of transition decreasing the energetic barrier.

Keywords: DPG, thermal decomposition, solvent.

Introduction

In the last years, systematic studies have been performed to study the kinetics and the mechanism of thermal decomposition reaction of 1, 2, 4, 5-tetroxanes, cyclic diperoxides which today have an important role in oxidation reactions of organic compounds with biological interest. [1-3]. These kinetic studies are intended to analyze the effect of substituents and solvents on the parameters governing the rate of reaction Steric, inductive, mesomeric and stereoelectronic effects of the different substituents on peroxidic ring are considered in relation by the force of peroxidic link (energy of

break of the link brought into play), which is debilitated in the initial step unimolecular of thermolysis. The general mechanism in solution belongs applicable to all the members of the family of them 1,2,4,5-tetroxanos, the unimolecular thermal decomposition can take place by two different kinds: a) a stepwise homolysis initiated by one peroxidic bond rupture with a birradical as intermediate. b) a concerted type of mechanism where bond breaking and bond making may occur simultaneously. However, for both processes in the thermolysis of tetroxanes give the corresponding ketone or aldehyde and molecular oxygen as reaction products. 1,2,4,5-tetroxanes shows the most significant solvent effect, in this work its thermal decomposition of 3,6-dibutanal-1,2,4,5-tetroxano (DPG, figure 1) in methanol solution have been studied examining the influence of this relatively polar medium.DPG presents a beak of absorbance to low lengths of wave presenting stable readings and fulfilling of Lambert Beer's Law to 204 nm, up to concentrations of the order of 10^{-6} M [4].In this paper a study of the thermal decomposition of this cyclic diperoxide in methanol is approached by UV spectroscopy, as an alternative methodology for the determination of kinetic parameters governing the thermolysis reaction of such compounds.



Figure 1. 3,6-dibutanal-1,2,4,5-tetroxane (DPG)

Materials and Methods

Purification of Solvents

Methanol (Merck, p.a.) was purified by fractionated distillation from disodium salt of ethylenediaminetetraacetic acid (disodium EDTA) in order to eliminate traces of metal ions that could be present in the commercial solvent [7]. GC analysis did not reveal the presence of organic impurities.

Synthesis Diperoxide

DPG was prepared by dropwise addition of glutaraldehyde in ethanol solution to a vigorously stirred and cooled (-20°C) mixture of 60% hydrogen peroxide and sulfuric acid (18M). After stirring for 2h more at the same temperature, the microcrystalline solid obtained was repeatedly washed with water and further purified by recrystallizing from methanol until a constant melting point was attained (192°C). The DPG purity was checked by GC in methanol solution.

Kinetic Methods

Pyrex glass tubes (4mm i.d., 70 mm length) filled with the appropriate volume of DPG in methanol solution, with added n-octane (internal standard) were thoroughly degassed in the vacuum line at -196°C and then sealed with a flame torch. These ampules were submerged in a thermostated silicone oil bath ($+0,1^{\circ}\text{C}$) to different temperatures and withdrawn after selected times, stopping the reaction cooling rapidly to 0°C . The closed ampules were conserved to -18°C up to his analysis for UV spectroscopy.

Analysis of Remaining Reagent.

The remaining DPG was performed by UV spectrophotometer; it marks CamSpec M330 at 204 nm.

Calculation Methods

The first order rate constant values were calculated by a least mean square data treatment ($r > 0,995$) and the activation parameters values of the thermal decomposition obtained from computational method employing the Eyring and Arrhenius equation parameters.

Results and Discussion

The thermal decomposition reaction of DPG in methanol solution in the range of temperature from $130,0$ to $166,0^{\circ}\text{C}$ and initial concentration $1,55 \times 10^{-3} \text{ mol L}^{-1}$ follow first order kinetic laws up to ca. 60 % diperoxide conversions (Fig. 2). In Table 1 we present values of the rate constants of reaction depending on the temperature. The above mentioned values are bigger than the obtained ones for other tetroxanes substituted in the same solvent (Table 2), which indicates that the substituent stabilizes the transition state by lowering the energy barrier.

The temperature influence on the rate constants values corresponding to the unimolecular reaction studied can be represented by the following Arrhenius equation

$$\ln k \text{ (s}^{-1}\text{)} = (16.7013 \pm 1.0) - (9745.43 \pm 700) / R T$$

The linearity of that equation ($r = 0,997$), in a wide range of temperatures (36°C) suggests that his parameters belong to a single process, which could be its unimolecular cleavage of the O-O bond as the initial step which could be its unimolecular cleavage of the O-O bond as the initial bond breaking step, as also it is described for the thermolysis of other tetroxanes in methanol [8]. The value of the activation energy observed ($E_a = 20.0 \text{ kcal mol}^{-1}$) gives us an indication that the two mechanisms postulated to interpret the thermolysis of other cyclic diperoxides, the mechanism for step is the most probable and the reaction is initiated with the formation of an intermediate biradical.

Theoretical calculations performed via uBHandLYP method of the decomposition reaction in gas phase of 3,6-dimethyl-1,2,4,5-tetroxane (a disubstituted tetroxane) following both reaction paths in stages and concerted, respectively, confirm the fact that the mechanism takes place in stages with an activation energy equal to 31.6 kcal/mol which is lower than the corresponding activation energy for a concerted mechanism (i.e. 72.8 kcal/mol) [9].

The activation parameters ($\Delta H^{\ddagger} = 18.5 \pm 0.9 \text{ kcal mol}^{-1}$ and $\Delta S^{\ddagger} = -45.2 \pm 0.6 \text{ cal K}^{-1} \text{ mol}^{-1}$) were worked out from the Eyring equation. They are showed in the Table 2, where bibliographic data are included to be compared. The activation parameter values obtained for the thermal decomposition reaction of DPG in methanol and in other tetroxanes (Table 2) indicate nearly ΔG^{\ddagger} values in the same studied solvent.

The very negative entropy value for the DPG thermolysis in methanol solution reflects the decrease in degree of freedoms of the DPG molecules that take place when they pass to a rather more rigid transition state, where the rupture of an O-O linkage might be assisted by the methanol solvent molecules. We believe that the interaction between the molecule and the solvent diperoxide takes place in the initial stage of the thermal decomposition reaction.

Conclusions

The thermal decomposition of the DPG in methanol, in the area of temperatures $130 - 166^{\circ}\text{C}$ and concentrations of the order of $1,55 \times 10^{-3} \text{ mol L}^{-1}$, fulfilled satisfactorily a kinetic law of the first order up to conversions of c.a. 60 % of diperoxide. Under experimental conditions, the values of activation parameters at initial stage of the reaction ($\Delta H^{\ddagger} = 18.5 \pm 0.9 \text{ kcal mol}^{-1}$, $\Delta S^{\ddagger} = -45.2 \pm 0.6 \text{ cal K}^{-1} \text{ mol}^{-1}$ and $19,36 \text{ kcal mol}^{-1}$) indicate us a mechanism initiated by the homolytic break of the peroxidic link for DPG that leads to the formation of an intermediary of the thermolysis (biradical), and this is determinant stage for the reaction. The theoretical calculations also support this mechanism.

The reaction products and the activation parameter values contribute to postulate the mechanism for the thermolysis of the DPG in methanol solution. The thermolysis would occur through a well known mechanism of decomposition already advanced for the tetroxanes, which begins with the homolytic

rupture of the peroxydic bond leading to the formation of an intermediate birradical and then C-O bond ruptures giving two molecules of glutaraldehyde and one oxygen molecule as final products. The large and negative activation entropy value (-45.2 cal / mol K) suggests that it makes up an interaction between solvent molecules and DPG.

References

1. Davies AG, Organic Peroxides, Butterworths, London. 1961.
2. Yamaguchi K, Takada K, Otsuji Y and Mizuno K, Organic Peroxides, Cap.1, 2-97. John Wiley & Sons Ltd. 1992.
3. Lawley KP, Advances in Chemical Physics, Vol. 69 I and II, Wiley, New York. 1987.
4. Wilson T, Landis ME, Baumstark AL and Bartlett PD. Solvent effects on the chemiluminescent decomposition of tetramethyl-1,2-dioxetane. Competitive dark pathways. J. Am. Chem. Soc. 1973;95: 4765-4766.
5. Kapstein R, Fráter-Schröder M, Oosterhoff LJ. Chemically induced dynamic nuclear polarization: NMR enhancements in biradical products. Chem. Phys. Letter. 1971; 12: 16-19.
6. Schuster GB, Schmidt SP. Chemiluminescence of Organic Compounds. Adv. Phys. Org. Chem. 1982;18: 187-238.
7. Weissberger, Proskauer, Riddick, Toops, Technique of Organic Chemistry. Vol VII. Organic Solvents. Physical Properties and Methods of Purification. 2nd ed. Interscience 1965.
8. Reguera M, Frette S, Romero JM, Jorge NL, Castro EA. Synthesis and thermal decomposition reaction of 3,6-dibutanoic-1,2,4,5-tetroxane in solution. BenSci. 2012; 4:1-4
9. Bordón AG, Profeta MI, Romero JM, Jorge NL, Hernandez-Laguna A and Grand AA. Thermolysis reaction of 3,6-dimethyl-1,2,4,5-tetroxane(DMT). WATOC2014.
- 10.

Table 1. First order rate constant values for DPG thermal decomposition reaction in methanol solution at different temperatures.

Temperature/°C	$k \times 10^4 \text{ s}^{-1}$
130	6.02
140	8.72
150	19.67
166	40.27

Table 2. First order rate constant values for tetroxanes substituted thermal decomposition reaction in methanol solution at 140°C

Tetroxane	$k \times 10^6 \text{ s}^{-1}$
DPAG ^a	6.9
ACDP ^b	14.0
DFT ^c	136.0
DPG	872.0

^aDPAG: Glutaraldehydeacidcyclicdiperoxide

^bACDP: Acetonecyclicdiperoxide

^cDFT: Benzaldehyde cyclic diperoxide




Figure 2. Kinetic of DPG thermal decomposition reaction ($1.55 \times 10^{-3} \text{ mol L}^{-1}$) in methanol solution at different temperatures

Table 2. Activation Parameters for DPAG thermolysis in methanol solution for different tetroxanes.

Tetroxane	ΔH^\ddagger Kcal mol ⁻¹	ΔS^\ddagger cal K ⁻¹ mol ⁻¹	ΔG^\ddagger Kcal mol ⁻¹
DPAG	25.2±1.0	-21.8±2.6	34.4±1.05
ACDP	26.9±0.3	-16.9±0.6	34.0±0.3
DFT	24,8±0.8	-16.7±1.8	31.8±0.8
DPG	18.5±0.9	-45.2±0.6	37.6±0.9




Figure 3.Eyring plot corresponding to the thermal decomposition reaction of DPG in methanol.