

Fractional  
Differential  
Calculus

*With Compliments  
of the Author*

Zagreb, Croatia

Volume 12, Number 2, December 2022

*Bahtiyar Bayraktar\* and Juan E. Nápoles Valdés*

*Hermite-Hadamard weighted integral inequalities  
for  $(h, m)$ -convex modified functions*

# FRACTIONAL DIFFERENTIAL CALCULUS

## AIMS AND SCOPE

'Fractional Differential Calculus' ('FDC') aims to publish original research papers on fractional differential and integral calculus, fractional differential equations and related topics. Specifically, contributions on both the mathematical and the numerical analysis of fractional differential calculus in engineering and sciences are welcome. Only papers of the highest quality will be accepted for publication.

'FDC' is published semiannually, in June and December.

## SUBMISSION

Manuscripts should be submitted electronically in Adobe PDF format to the 'FDC' Editorial Office through the provided web interface.

The submitter of the paper to the website is referred to as the corresponding author and is the one expected to manage all communication and correspondence with 'FDC' regarding the paper. Authors may indicate their preference with regard to which member of the Editorial Board they consider appropriate for the article. However, assignment to that particular member is not assured.

Manuscripts are accepted for refereeing on the understanding that the same work has not been published (except in the form of an abstract), and is not under consideration for publication elsewhere.

In order to facilitate refereeing, copies of those papers (whether by the author or someone else) which are essential and referred to in the manuscript but are not conveniently accessible, should be enclosed.

The publisher strongly encourages submission of manuscripts written in TeX or one of its variants LaTeX, AMSTeX or AMSLaTeX. On acceptance of the paper, authors will be asked to send the file by electronic mail. The file must be the one from which the accompanying manuscript (finalized version) was printed out. A PDF file of this final version must also be enclosed.

## PREPARATION OF MANUSCRIPT

The following data should be included on title page: the title of the article, author's name (no degrees) author's affiliation, e-mail addresses, mailing address of the corresponding author, and running head less than 60 characters.

### Abstract, Keywords, Subject classification

The manuscript must be accompanied by a brief abstract, no longer than 100-150 words. It should make minimal use of mathematical symbols and displayed formulas. A list of 4-5 key words must be given. Mathematics Subject Classification (2010) with primary (and secondary) subject classification codes must be included.

### Figures

Figures should be prepared in a digital form suitable for direct reproduction, grey scale only, preferably at resolution of 300 dpi or higher, sized to fit a journal page, and in EPS, TIFF or JPEG format. Figures should have a short descriptive title and should be numbered consecutively in order of appearance in the text. The position of each figure in the text should be indicated.

### References

Bibliographic references should be listed alphabetically at the end of the article. The authors should consult Mathematical Reviews for the standard abbreviations of journal names.

### PROOFS

Galley proofs will be sent in PDF format to corresponding author. It should be returned in a few days.

### PAGE CHARGES, OFFPRINTS

There are no page charges. Authors will receive PDF file of the printed article free of charge. Additional offprints may be ordered from FDC prior to publication.

FDC is published by Publishing House **ELEMENT**, Zagreb, Croatia.

All correspondence and subscription orders should be addressed to the Editorial Office:

[www.ele-math.com](http://www.ele-math.com)

e-mail: [fdc@ele-math.com](mailto:fdc@ele-math.com)

Fractional Differential Calculus  
Editorial Office  
Menceticeva 2, 10000 Zagreb, Croatia  
Fax: +385 1 6008799

## HERMITE–HADAMARD WEIGHTED INTEGRAL INEQUALITIES FOR $(h, m)$ –CONVEX MODIFIED FUNCTIONS

BAHTIYAR BAYRAKTAR\* AND JUAN E. NÁPOLES VALDÉS

(Communicated by S. S. Dragomir)

*Abstract.* In this paper, some new integral inequalities of the Hermite–Hadamard type are were obtained for  $(h, m)$ -convex modified functions. The results are obtained on the basis of the introduced definition of a generalized weighted integral operator by using the convexity property, the well-known Hölder's inequality and its modification. Some results existing in the literature are some special cases of our results.

### 1. Introduction

A function  $\psi : [\rho, \sigma] \rightarrow \mathbb{R}$ , is said to be convex if  $\psi(\theta x_1 + (1 - \theta)x_2) \leq \theta\psi(x_1) + (1 - \theta)\psi(x_2)$  holds  $\forall x_1, x_2 \in [\rho, \sigma]$  and  $\theta \in [0, 1]$ . If the above inequality is reversed, then the function  $\psi$  will be concave on  $[\rho, \sigma]$ .

Convex functions have been summed up widely hypothetically ; these extensions incorporate the  $m$ -convex,  $r$ -concave,  $h$ -convex,  $s$ -convex,  $(h, m)$ -convex functions and numerous others. Readers interested in its multiple ramifications and extensions can consult [26], where a fairly complete overview of the development of the convex function concept was presented.

**DEFINITION 1.** [32] Let  $\psi : [0, \sigma] \rightarrow \mathbb{R}$  and  $m \in [0, 1]$ , If  $\forall x_1, x_2 \in [0, \sigma]$  and  $\theta \in [0, 1]$  the inequality

$$\psi(\theta x_1 + (1 - \theta)x_2) \leq \theta\psi(x_1) + m(1 - \theta)\psi(x_2) \quad (1)$$

holds. Then  $\psi$  is said to be  $m$ -convex on  $[0, \sigma]$ .

If the above inequality is holds in reverse, then we say that the function  $\psi$  is  $m$ -concave.

The following definitions are the successive extensions of the concept of convex function and, as we will see later, they are the particular cases of our Definition.

*Mathematics subject classification* (2020): 26D10, 26A51, 26A33.

*Keywords and phrases:* Hermite-Hadamard type integral inequality, Hölder's integral inequality, weighted integral operators,  $(h, m)$ -convex modified functions.

\* Corresponding author.

DEFINITION 2. [9, 15] Let  $\psi : [0, \sigma] \rightarrow \mathbb{R}$  and  $s \in (0, 1]$ . If,  $\forall x_1, x_2 \in [0, \sigma]$  and  $\theta \in [0, 1]$  the inequality  $\psi(\theta x_1 + (1 - \theta)x_2) \leq \theta^s \psi(x_1) + (1 - \theta^s) \psi(x_2)$  is valid, then on  $[0, \sigma]$  the function  $\psi$  is called the  $s$ -convex(in the first sense).

DEFINITION 3. [9, 15] Let  $\psi : [0, \sigma] \rightarrow \mathbb{R}$  and  $s \in (0, 1]$ . If  $\forall x_1, x_2 \in [0, \sigma]$  and  $\theta \in [0, 1]$  the inequality  $\psi(\theta x_1 + (1 - \theta)x_2) \leq \theta^s \psi(x_1) + (1 - \theta)^s \psi(x_2)$  is valid, then on  $[0, \sigma]$  the function  $\psi$  is called the  $s$ -convex(in the second sense).

DEFINITION 4. [34] Let  $\psi : [0, \sigma] \rightarrow \mathbb{R}$  and  $s \in [-1, 1]$ . If  $\forall x_1, x_2 \in [0, \sigma]$  and  $\theta \in (0, 1)$  the inequality  $\psi(\theta x_1 + (1 - \theta)x_2) \leq \theta^s \psi(x_1) + (1 - \theta)^s \psi(x_2)$  is valid, then on  $[0, \sigma]$  the function  $\psi$  is called the extended  $s$ -convex.

In [21], the present class of  $(a, m)$ -convex functions as follows.

DEFINITION 5. Let  $\psi : [0, \sigma] \rightarrow \mathbb{R}$  and  $\alpha, m \in [0, 1]$ . If  $\forall x_1, x_2 \in [0, \sigma]$  and  $\theta \in [0, 1]$  and the inequality  $\psi(\theta x_1 + m(1 - \theta)x_2) \leq \theta^\alpha \psi(x_1) + m(1 - \theta^\alpha) \psi(x_2)$  is valid, then on  $[0, \sigma]$  the function  $\psi$  is called the  $(a, m)$ -convex.

In [20], the following definition is introduced.

DEFINITION 6. Let  $h : [0, 1] \rightarrow \mathbb{R}$  and  $\psi : [0, \sigma] \rightarrow [0, +\infty)$  are non-negative functions and the function  $h$  is not identically zero. If  $\forall x_1, x_2 \in [0, \sigma]$  and  $\theta \in [0, 1]$  the inequality

$$\psi(\theta x_1 + m(1 - \theta)x_2) \leq h(\theta) \psi(x_1) + mh(1 - \theta) \psi(x_2) \quad (2)$$

is fulfilled for  $m \in [0, 1]$ , then on  $[0, \sigma]$  the function  $\psi$  is called the  $(h, m)$ -convex.

If the above inequality is reversed, then  $\psi$  is said to be  $(h, m)$ -concave. Note that in Definition 6, if we take  $h(\theta) = \theta$ , then we get the definition of an  $m$ -convex function, and if, additionally, we take  $m = 1$ , then we get the definition of classical convexity.

In [25], the authors presented the class of  $s - (a, m)$ -convex functions as follows (“redefined” in [35]).

DEFINITION 7. Let  $\psi : [0, +\infty) \rightarrow [0, +\infty)$  and  $(a, m) \in [0, 1]^2$ , and  $s \in (0, 1]$ . If,  $\forall \rho, \sigma \in [0, +\infty)$  and  $\theta \in [0, 1]$  the inequality

$$\psi(\theta \rho + m(1 - \theta)\sigma) \leq \theta^{as} \psi(\rho) + m(1 - \theta^{as}) \psi(\sigma)$$

is valid, then the function  $\psi$  is called the  $s - (a, m)$ -convex in the first sense.

DEFINITION 8. Let  $\psi : [0, +\infty) \rightarrow [0, +\infty)$  and  $a, m \in [0, 1]$ , and  $s \in (0, 1]$ . If,  $\forall \rho, \sigma \in [0, +\infty)$  and  $\theta \in [0, 1]$  the inequality

$$\psi(\theta \rho + m(1 - \theta)\sigma) \leq (\theta^a)^s \psi(\rho) + m(1 - \theta^a)^s \psi(\sigma),$$

is valid, then the function  $\psi$  is called the  $s - (a, m)$ -convex in the second sense.

On the basis of these definitions, we will present the classes of functions that will be the basis of our work ([3]).

DEFINITION 9. Let  $h : [0, 1] \rightarrow [0, 1]$  and  $\psi : [0, \sigma] \rightarrow [0, +\infty)$  be non-negative functions and the function  $h$  is not identically zero. If,  $\forall x_1, x_2 \in [0, +\infty)$  and  $\theta \in [0, 1]$  the inequality

$$\psi(\theta x_1 + m(1 - \theta)x_2) \leq h^s(\theta)\psi(x_1) + m(1 - h^s(\theta))\psi(x_2) \quad (3)$$

is valid for  $m \in [0, 1]$  and  $s \in [-1, 1]$ , then on  $[0, +\infty)$  the function  $\psi$  is called the  $(h, m)$ -convex modified of the first type.

DEFINITION 10. Let  $h : [0, 1] \rightarrow [0, 1]$  and  $\psi : [0, \sigma] \rightarrow [0, +\infty)$  be non-negative functions and the function  $h$  is not identically zero. If,  $\forall x_1, x_2 \in [0, +\infty)$  and  $\theta \in [0, 1]$  the inequality

$$\psi(\theta x_1 + m(1 - \theta)x_2) \leq h^s(\theta)\psi(x_1) + m(1 - h(\theta))^s\psi(x_2) \quad (4)$$

is valid for  $m \in [0, 1]$  and  $s \in [-1, 1]$ , then on  $[0, +\infty)$  the function  $\psi$  is called the  $(h, m)$ -convex modified of the second type.

REMARK 1. From Definitions 9 and 10, we have

1. If  $h(\theta) = \theta^a$  with  $a \in (0, 1]$ , then  $\psi$  is a  $s - (a, m)$ -convex function on  $[0, +\infty)$ .
2. If  $s = 1$ , then  $\psi$  is an  $(h, m)$ -convex function on  $[0, +\infty)$ .
3. If  $h(\theta) = \theta$ ,  $s \in (0, 1]$  and  $m = 1$ , then  $\psi$  is a  $s$ -convex function on  $[0, +\infty)$ .
4. If  $h(\theta) = \theta$ ,  $s \in [-1, 1]$  and  $m = 1$ , then  $\psi$  is an extended  $s$ -convex function on  $[0, +\infty)$ .
5. If  $h(\theta) = \theta$  and  $s = m = 1$ , then  $\psi$  is a convex function on  $[0, +\infty)$ .

One of the most important inequalities, that has attracted the attention many inequality experts in the last few decades, is the famous Hermite–Hadamard inequality:

$$\psi\left(\frac{\rho + \sigma}{2}\right) \leq \frac{1}{\sigma - \rho} \int_{\rho}^{\sigma} \psi(\theta) d\theta \leq \frac{\psi(\rho) + \psi(\sigma)}{2} \quad (5)$$

which holds for any function  $\psi$  convex on the interval  $[\rho, \sigma]$ . This inequality was proved by the French mathematicians Charles Hermite in 1883 and independently, 10 years later, by Jacques Hadamard. The importance of this inequality is that it makes it possible to estimate the mean value of a convex function, moreover, it provides a refinement of the well-known Jensen inequality. Several results can be consulted in [1, 8, 11, 4, 7, 12, 13, 14, 16, 18, 20, 23, 22, 27] as well as the references therein for more information and other extensions of the Hermite–Hadamard inequality.

To promote the understanding of the subject, we firstly presented the definition of the classical Riemann–Liouville fractional integrals (with  $0 \leq \rho < \theta < \sigma \leq \infty$ ).

DEFINITION 11. Let  $\psi \in L_1[\rho, \sigma]$ . Then, the Riemann-Liouville fractional integrals of order  $\alpha \in \mathbb{C}$ ,  $\Re(\alpha) > 0$  are defined by (right and left respectively):

$$\begin{aligned} {}^\alpha I_{\rho+} \psi(x_1) &= \frac{1}{\Gamma(\alpha)} \int_{\rho}^{x_1} (x_1 - \theta)^{\alpha-1} \psi(\theta) d\theta, \quad x_1 > \rho, \\ {}^\alpha I_{\sigma-} \psi(x_1) &= \frac{1}{\Gamma(\alpha)} \int_{x_1}^{\sigma} (\theta - x_1)^{\alpha-1} \psi(\theta) d\theta, \quad x_1 < \sigma, \end{aligned}$$

where  $\Gamma$  is the Euler gamma function.

The basis of our work is the next definition of the weighted integral operators:

DEFINITION 12. Let  $\psi \in L_1[\rho, \sigma]$  and  $w : [0, 1] \rightarrow \mathbb{R}^+$ ,  $w \in C([0, 1])$  having the first and second-order derivatives piecewise continuous on  $[0, 1]$ ,  $w(0) = 0$  and  $w(1) = 0$ . Then the weighted fractional integrals are defined by (right and left respectively):

$$\begin{aligned} I_{\rho+} \psi(x_1) &= \int_{\rho}^{x_1} w'' \left( \frac{x_1 - \theta}{\frac{\rho + \sigma}{2} - \rho} \right) \psi(\theta) d\theta, \quad x_1 > \rho \\ I_{\sigma-} \psi(x_1) &= \int_{x_1}^{\sigma} w'' \left( \frac{\theta - x_1}{\sigma - \frac{\rho + \sigma}{2}} \right) \psi(\theta) d\theta, \quad x_1 < \sigma. \end{aligned}$$

REMARK 2. If  $w''(\theta) = \frac{(\frac{\sigma-\rho}{2})^{\alpha-1} \theta^{\alpha-1}}{\Gamma(\alpha)}$ , then we obtain the Riemann-Liouville fractional integral, right and left.

REMARK 3. By putting  $w''(\theta) \equiv 1$ , we obtain the classical Riemann integral.

In this paper, within the framework of generalized integral operators (Definition 12), we presented some variants of the inequality (5) for  $(h, m)$ -convex modified functions.

## 2. Hermite–Hadamard type inequalities for $(h, m)$ -convex modified functions

To establish our results, we need the following Lemma.

LEMMA 1. Let  $\psi : I = [\rho, \sigma] \rightarrow \mathbb{R}$  and  $\psi \in C^2(I^\circ)$ . If  $\psi'' \in L_1(I)$ , then

$$\begin{aligned} & -\frac{1}{2} \left[ w'(1)\psi(\rho) - w'(0)\psi\left(\frac{\rho + \sigma}{2}\right) + w'(1)\psi\left(\frac{\rho + \sigma}{2}\right) - w'(0)\psi(\sigma) \right] \quad (6) \\ & + \frac{1}{\sigma - \rho} \left( I_{\rho+} \psi\left(\frac{\rho + \sigma}{2}\right) + I_{\sigma-} \psi\left(\frac{\rho + \sigma}{2}\right) \right) \\ & = \frac{(\sigma - \rho)^2}{8} \int_0^1 w(\theta) \left[ \psi''\left(\theta\rho + (1-\theta)\frac{\rho + \sigma}{2}\right) + \psi''\left(\theta\frac{\rho + \sigma}{2} + (1-\theta)\sigma\right) \right] d\theta. \end{aligned}$$

*Proof.* Denoting

$$\begin{aligned} I &= \int_0^1 w(\theta) \left[ \psi'' \left( \theta\rho + (1-\theta) \frac{\rho+\sigma}{2} \right) + \psi'' \left( \theta \frac{\rho+\sigma}{2} + (1-\theta)\sigma \right) \right] d\theta \\ &= I_1 + I_2. \end{aligned}$$

If we integrate by parts and we take into account the definition of  $w(\theta)$ , we obtain

$$\begin{aligned} I_1 &= \int_0^1 w(\theta) \psi'' \left( \theta\rho + (1-\theta) \frac{\rho+\sigma}{2} \right) d\theta \\ &= \frac{2}{\sigma-\rho} \int_0^1 w'(\theta) \psi' \left( \theta\rho + (1-\theta) \frac{\rho+\sigma}{2} \right) d\theta \\ &= \frac{2}{\sigma-\rho} \left[ \frac{2w'(\theta)}{\rho-\sigma} \psi \left( \theta\rho + (1-\theta) \frac{\rho+\sigma}{2} \right) \Big|_0^1 \right. \\ &\quad \left. - \frac{2}{\rho-\sigma} \int_0^1 w''(\theta) \psi \left( \theta\rho + (1-\theta) \frac{\rho+\sigma}{2} \right) d\theta \right] \\ &= \frac{-4}{(\sigma-\rho)^2} \left[ \left( w'(1)\psi(\rho) - w'(0)\psi \left( \frac{\rho+\sigma}{2} \right) \right) \right. \\ &\quad \left. - \int_0^1 w''(\theta) \psi \left( \theta\rho + (1-\theta) \frac{\rho+\sigma}{2} \right) d\theta \right] \end{aligned}$$

By using the change of the variable  $z = \theta\rho + (1-\theta) \frac{\rho+\sigma}{2}$  for  $\theta \in [0, 1]$ , which gives

$$I_1 = \frac{-4}{(\sigma-\rho)^2} \left[ w'(1)\psi(\rho) - w'(0)\psi \left( \frac{\rho+\sigma}{2} \right) \right] + \frac{8}{(\sigma-\rho)^3} I_{\rho+\psi} \left( \frac{\rho+\sigma}{2} \right).$$

Similarly, we obtain

$$I_2 = \frac{-4}{(\sigma-\rho)^2} \left[ w'(1)\psi \left( \frac{\rho+\sigma}{2} \right) - w'(0)\psi(\sigma) \right] + \frac{8}{(\sigma-\rho)^3} I_{\sigma-\psi} \left( \frac{\rho+\sigma}{2} \right).$$

Then,

$$\begin{aligned} I &= I_1 + I_2 \\ &= \frac{-4}{(\sigma-\rho)^2} \left[ w'(1)\psi(\rho) - w'(0)\psi \left( \frac{\rho+\sigma}{2} \right) + w'(1)\psi \left( \frac{\rho+\sigma}{2} \right) - w'(0)\psi(\sigma) \right] \\ &\quad + \frac{8}{(\sigma-\rho)^3} \left( I_{\rho+\psi} \left( \frac{\rho+\sigma}{2} \right) + I_{\sigma-\psi} \left( \frac{\rho+\sigma}{2} \right) \right). \end{aligned}$$

From here, it is very easy to obtain the required equality.  $\square$

REMARK 4. If we use

$$w(\theta) = \begin{cases} \theta^\alpha, & \theta \in [0, \frac{1}{2}) \\ (1 - \theta)^\alpha, & \theta \in [\frac{1}{2}, 1] \end{cases},$$

then

1. For  $\alpha > 1$ , from the lemma above with only  $\psi(\theta\rho + m(1 - \theta)\sigma)$ , the Lemma 2.1 of [5] is obtained;
2. If we choose  $\alpha = 1$  and, from the lemma above with only  $\psi(\theta\rho + m(1 - \theta)\sigma)$ , the Lemma 2.1 of [6] is obtained;
3. If we choose  $\alpha = 2$  and consider the functions  $\psi(\theta\rho + (1 - \theta)\sigma)$  and  $\psi(\theta\sigma + (1 - \theta)\rho)$ , then, from (6), we obtain Lemma 2 from [31].

REMARK 5. If we consider the functions  $\psi(\theta\rho + (1 - \theta)\sigma)$  and  $\psi(\theta\sigma + (1 - \theta)\rho)$  use  $w(\theta) = \theta(1 - \theta)^\alpha$ , then, from (6), we obtain Lemma 3.1 from [5].

REMARK 6. Under the same requirements of the previous Remark, and if we consider  $w(\theta) = \frac{1 - (1 - \theta)^{\alpha+1} - \theta^{\alpha+1}}{\alpha+1}$ , from the Lemma above with only  $\psi(\theta\rho + (1 - \theta)\sigma)$ , the Lemma 2.1 of [33] is obtained.

REMARK 7. If we put  $w(\theta) = \theta(1 - \theta)$ , then from (6), we obtain Lemma 2.1 of [35]. However, if we put  $w(\theta) = \frac{\theta^{(\alpha+1)}}{\alpha(\alpha+1)\Gamma(\alpha)}$ , we obtain the following new equality for Fractional Integral of Riemann-Liouville:

$$\begin{aligned} & -\frac{1}{2\Gamma(\alpha+1)} \left[ \psi(\rho) + \psi\left(\frac{\rho+\sigma}{2}\right) \right] - \frac{\sigma-\rho}{4\Gamma(\alpha+2)} \left[ \psi'(\rho) + \psi'\left(\frac{\rho+\sigma}{2}\right) \right] \quad (7) \\ & + \frac{2^{\alpha-1}}{(\sigma-\rho)^\alpha} \left[ {}^\alpha I_{\rho+} \psi\left(\frac{\rho+\sigma}{2}\right) + {}^\alpha I_{\sigma-} \psi\left(\frac{\rho+\sigma}{2}\right) \right] \\ & = \frac{(\sigma-\rho)^2}{8\Gamma(\alpha+2)} \int_0^1 \theta^{\alpha+1} \left[ \psi''\left(\theta\rho + (1-\theta)\frac{\rho+\sigma}{2}\right) + \psi''\left(\theta\frac{\rho+\sigma}{2} + (1-\theta)\sigma\right) \right] d\theta. \end{aligned}$$

THEOREM 1. Let  $\psi : I = [\rho, \sigma] \rightarrow \mathbb{R}$  and  $\psi \in C^2(I^\circ)$  with  $\psi'' \in L_1(I)$ . If  $|\psi''|$  is a convex function ( $h(\theta) = \theta$  and  $s = m = 1$ ), then the inequality

$$\begin{aligned} & \left| \frac{2^{\alpha-1}}{(\sigma-\rho)^\alpha} \left[ {}^\alpha I_{\rho+} \psi\left(\frac{\rho+\sigma}{2}\right) + {}^\alpha I_{\sigma-} \psi\left(\frac{\rho+\sigma}{2}\right) \right] - \frac{\psi(\rho) + \psi\left(\frac{\rho+\sigma}{2}\right)}{2\Gamma(\alpha+1)} \right. \\ & \quad \left. - \frac{(\sigma-\rho) \left[ \psi'(\rho) + \psi'\left(\frac{\rho+\sigma}{2}\right) \right]}{4\Gamma(\alpha+2)} \right| \quad (8) \\ & \leq \frac{(\sigma-\rho)^2}{8\Gamma(\alpha+2)} \left[ \frac{|\psi''(\rho)| - |\psi''(\sigma)|}{\alpha+3} + \frac{|\psi''\left(\frac{\rho+\sigma}{2}\right)| + |\psi''(\sigma)|}{\alpha+2} \right]. \end{aligned}$$

holds  $\forall \alpha > 0$ .

*Proof.* It is easy to see that for the right-hand side of the identity (7), by using the properties of the modulus and taking into account the convexity of the function  $\psi$ , we obtain:

$$\begin{aligned} & \left| \int_0^1 \theta^{\alpha+1} \left[ \psi'' \left( \theta \rho + (1-\theta) \frac{\rho+\sigma}{2} \right) + \psi'' \left( \theta \frac{\rho+\sigma}{2} + (1-\theta) \sigma \right) \right] d\theta \right| \\ & \leq \frac{|\psi''(\rho)| - |\psi''(\sigma)|}{\alpha+3} + \frac{|\psi''(\frac{\rho+\sigma}{2})| + |\psi''(\sigma)|}{\alpha+2}. \end{aligned}$$

This inequality shows the validity of (8).  $\square$

Now we get some new inequalities of the Hermite-Hadamard type.

**THEOREM 2.** Let  $\psi : I = [\rho, \frac{\sigma}{m}] \rightarrow \mathbb{R}$  and  $\psi \in C^2(I^\circ)$  with  $\psi'' \in L_1(I)$ . If  $|\psi''|$  is  $(h, m)$ -convex modified of the second sense on  $I$ , then

$$\begin{aligned} & \left| R_w + \frac{1}{\sigma - \rho} \left( I_{\rho+} \psi \left( \frac{\rho + \sigma}{2} \right) + I_{\sigma-} \psi \left( \frac{\rho + \sigma}{2} \right) \right) \right| \tag{9} \\ & \leq \frac{(\sigma - \rho)^2}{8} \left[ \left( |\psi''(\rho)| + \left| \psi'' \left( \frac{\rho + \sigma}{2} \right) \right| \right) \mathbb{A} + m \left( \left| \psi'' \left( \frac{\rho + \sigma}{2m} \right) \right| + \left| \psi'' \left( \frac{\sigma}{m} \right) \right| \right) \mathbb{B} \right], \end{aligned}$$

with

$$\begin{aligned} R_w &= -\frac{1}{2} \left[ w'(1) \psi(\rho) - w't(0) \psi \left( \frac{\rho + \sigma}{2} \right) + w'(1) \psi \left( \frac{\rho + \sigma}{2} \right) - w'(0) \psi(\sigma) \right], \\ \mathbb{A} &= \int_0^1 |w(\theta)| h^s(\theta) d\theta \quad \text{and} \quad \mathbb{B} = \int_0^1 |w(\theta)| (1 - h(\theta))^s d\theta. \end{aligned}$$

*Proof.* From Lemma 1, it follows that

$$\begin{aligned} & \left| R_w + \frac{1}{(\sigma - \rho)} \left( I_{\rho+} \psi \left( \frac{\rho + \sigma}{2} \right) + I_{\sigma-} \psi \left( \frac{\rho + \sigma}{2} \right) \right) \right| \\ & \leq \frac{(\sigma - \rho)^2}{8} (I_1 + I_2). \end{aligned}$$

with

$$\begin{aligned} I_1 &= \int_0^1 |w(\theta)| \left| \psi'' \left( \theta \rho + (1-\theta) \frac{\rho + \sigma}{2} \right) \right| d\theta \\ &= \int_0^1 |w(\theta)| \left| \psi'' \left( \theta \rho + m(1-\theta) \frac{\rho + \sigma}{2m} \right) \right| d\theta \end{aligned}$$

and

$$\begin{aligned} I_2 &= \int_0^1 |w(\theta)| \left| \psi'' \left( \theta \frac{\rho + \sigma}{2} + (1-\theta) \sigma \right) \right| d\theta \\ &= \int_0^1 |w(\theta)| \left| \psi'' \left( \theta \frac{\rho + \sigma}{2} + m(1-\theta) \frac{\sigma}{m} \right) \right| d\theta. \end{aligned}$$

To obtain the inequality (9), just use the  $(h, m)$ -convex modified of the second sense of  $|\psi''|$ , in  $I_1$  and  $I_2$ , apply the elementary properties of the module and group like terms.  $\square$

REMARK 8. For  $h(\theta) = \theta$ , that is, if  $\psi$  is a  $(s, m)$ -convex function, then

1. By taking into account Remark 4.1, it is not difficult to obtain Theorem 2.1 of [5];
2. By taking into account Remark 4.2, it is easy to obtain Theorem 2.1 of [6];
3. By taking into account Remark 5, we obtain Theorem 3.1 of [5].

REMARK 9. By taking into account Remark 4.3, it is easy to obtain Theorem 3 of [31] by considering that  $\psi$  that is a convex function that is, if  $h(\theta) = \theta$  and  $s = 1$ .

THEOREM 3. Let  $\psi : I = [\rho, \frac{\sigma}{m}] \rightarrow \mathbb{R}$  and  $\psi \in C^2(I^\circ)$  such that  $\psi'' \in L_1(I)$ . If  $|\psi|^q$  is  $(h, m)$ -convex modified of the second sense on  $I$ ,  $q \geq 1$ , then

$$\begin{aligned} & \left| R_w + \frac{1}{(\sigma - \rho)} \left( I_{\rho+} \psi \left( \frac{\rho + \sigma}{2} \right) + I_{\sigma-} \psi \left( \frac{\rho + \sigma}{2} \right) \right) \right| \\ & \leq \frac{(\sigma - \rho)^2}{8} \left( \int_0^1 |w(\theta)|^p dt \right)^{\frac{1}{p}} \\ & \quad \times \left\{ \left( |\psi''(\rho)|^q \mathbb{H}_1 + m \left| \psi'' \left( \frac{\rho + \sigma}{2m} \right) \right|^q \mathbb{H}_2 \right)^{\frac{1}{q}} \right. \\ & \quad \left. + \left( \left| \psi'' \left( \frac{\rho + \sigma}{2} \right) \right|^q \mathbb{H}_1 + m \left| \psi'' \left( \frac{\sigma}{m} \right) \right|^q \mathbb{H}_2 \right)^{\frac{1}{q}} \right\}, \end{aligned}$$

with

$$\begin{aligned} R_w &= -\frac{1}{2} \left[ w'(1)\psi(\rho) - w'(0)\psi \left( \frac{\rho + \sigma}{2} \right) + w'(1)\psi \left( \frac{\rho + \sigma}{2} \right) - w'(0)\psi(\sigma) \right], \\ \mathbb{H}_1 &= \int_0^1 h^s(\theta) dt \quad \text{and} \quad \mathbb{H}_2 = \int_0^1 (1 - h(\theta))^s dt. \end{aligned}$$

*Proof.* From Lemma 1 and using well-known Hölder’s integral inequality, we get

$$\begin{aligned} & \left| R_w + \frac{1}{(\sigma - \rho)} \left( I_{\rho+} \psi \left( \frac{\rho + \sigma}{2} \right) + I_{\sigma-} \psi \left( \frac{\rho + \sigma}{2} \right) \right) \right| \tag{10} \\ & \leq \frac{(\sigma - \rho)^2}{8} \left( \int_0^1 |w(\theta)|^p d\theta \right)^{\frac{1}{p}} \left\{ I_3^{\frac{1}{q}} + I_4^{\frac{1}{q}} \right\}, \end{aligned}$$

with  $I_3 = \int_0^1 |\psi''(\theta\rho + (1 - \theta)\frac{\rho + \sigma}{2})|^q d\theta$ ,  $I_4 = \int_0^1 |\psi''(\theta\frac{\rho + \sigma}{2} + (1 - \theta)\sigma)|^q d\theta$ .

Since  $|\psi''|^q$  is  $(h, m)$ -convex modified of the second sense on  $[\rho, \frac{\sigma}{m}]$ , we know that for  $\theta \in [0, 1]$

$$\left| \psi'' \left( \theta \rho + m(1-\theta) \frac{\rho + \sigma}{2m} \right) \right|^q \leq h^s(\theta) |\psi''(\rho)|^q + m(1-h(\theta))^s \left| \psi'' \left( \frac{\rho + \sigma}{2m} \right) \right|^q, \quad (11)$$

$$\left| \psi'' \left( \theta \frac{\rho + \sigma}{2} + m(1-\theta) \frac{\sigma}{m} \right) \right|^q \leq h^s(\theta) \left| \psi'' \left( \frac{\rho + \sigma}{2} \right) \right|^q + m(1-h(\theta))^s \left| \psi'' \left( \frac{\sigma}{m} \right) \right|^q \quad (12)$$

Hence, by substituting (11) and (12) in (10), we get

$$\begin{aligned} & \left( \int_0^1 |w(\theta)|^p d\theta \right)^{\frac{1}{p}} \left\{ I_3^q + I_4^q \right\} \leq \left( \int_0^1 |w(\theta)|^p d\theta \right)^{\frac{1}{p}} \\ & \times \left\{ \left( \int_0^1 \left[ h^s(\theta) |\psi''(\rho)|^q + m(1-h(\theta))^s \left| \psi'' \left( \frac{\rho + \sigma}{2m} \right) \right|^q \right] d\theta \right)^{\frac{1}{q}} \right. \\ & \left. + \left( \int_0^1 \left[ h^s(\theta) \left| \psi'' \left( \frac{\rho + \sigma}{2} \right) \right|^q + m(1-h(\theta))^s \left| \psi'' \left( \frac{\sigma}{m} \right) \right|^q \right] d\theta \right)^{\frac{1}{q}} \right\} \\ & = \left( \int_0^1 |w(\theta)|^p d\theta \right)^{\frac{1}{p}} \left\{ \left( |\psi''(\rho)|^q \mathbb{H}_1 + m \left| \psi'' \left( \frac{\rho + \sigma}{2m} \right) \right|^q \mathbb{H}_2 \right)^{\frac{1}{q}} \right. \\ & \left. + \left( \left| \psi'' \left( \frac{\rho + \sigma}{2} \right) \right|^q \mathbb{H}_1 + m \left| \psi'' \left( \frac{\sigma}{m} \right) \right|^q \mathbb{H}_2 \right)^{\frac{1}{q}} \right\} \end{aligned}$$

which completes the proof.  $\square$

REMARK 10. By considering  $w(\theta) = \theta(1-\theta)$  and taking  $\psi''$  as a function  $(\alpha, m)$ -convex, we obtain Theorem 3.1 of [35].

A variant of the previous result is given in the following theorem.

THEOREM 4. Let  $\psi : I = [\rho, \frac{\sigma}{m}] \rightarrow \mathbb{R}$  and  $\psi \in C^2(I^\circ)$  such that  $\psi'' \in L_1(I)$ . If  $|\psi|^q$  is  $(h, m)$ -convex modified of the second sense on  $I$ ,  $q > 1$ , then

$$\begin{aligned} & \left| R_w + \frac{1}{(\sigma - \rho)} \left( I_{\rho+} \psi \left( \frac{\rho + \sigma}{2} \right) + I_{\sigma-} \psi \left( \frac{\rho + \sigma}{2} \right) \right) \right| \quad (13) \\ & \leq \frac{(\sigma - \rho)^2}{8} \left( \int_0^1 |w(\theta)|^p d\theta \right)^{\frac{1}{p}} \\ & \times \left\{ \left( |\psi''(\rho)|^q \mathbb{H}_1 + m \left| \psi'' \left( \frac{\rho + \sigma}{2m} \right) \right|^q \mathbb{H}_2 \right)^{\frac{1}{q}} \right. \\ & \left. + \left( \left| \psi'' \left( \frac{\rho + \sigma}{2} \right) \right|^q \mathbb{H}_1 + m \left| \psi'' \left( \frac{\sigma}{m} \right) \right|^q \mathbb{H}_2 \right)^{\frac{1}{q}} \right\}, \end{aligned}$$

with

$$R_w = -\frac{1}{2} \left[ w'(1)\psi(\rho) - w'(0)\psi\left(\frac{\rho + \sigma}{2}\right) + w'(1)\psi\left(\frac{\rho + \sigma}{2}\right) - w'(0)\psi(\sigma) \right],$$

$$\mathbb{H}_1 = \int_0^1 |h^s(\theta)|d\theta \quad \text{and} \quad \mathbb{H}_2 = \int_0^1 (1 - h(\theta))^s d\theta.$$

*Proof.* From Lemma 1 and by using well-known power mean inequality, we have

$$\left| R_w + \frac{1}{(\sigma - \rho)} \left( I_{\rho+} \psi\left(\frac{\rho + \sigma}{2}\right) + I_{\sigma-} \psi\left(\frac{\rho + \sigma}{2}\right) \right) \right| \tag{14}$$

$$\leq \frac{(\sigma - \rho)^2}{8} \left( \int_0^1 |w(\theta)|d\theta \right)^{1-\frac{1}{q}} \left\{ I_5^{\frac{1}{q}} + I_6^{\frac{1}{q}} \right\},$$

with

$$I_5 = \left( \int_0^1 |w(\theta)| \left| \psi'' \left( \theta\rho + (1 - \theta)\frac{\rho + \sigma}{2} \right) \right|^q d\theta \right)$$

and

$$I_6 = \left( \int_0^1 |w(\theta)| \left| \psi'' \left( \theta\frac{\rho + \sigma}{2} + (1 - \theta)\sigma \right) \right|^q d\theta \right).$$

Since  $|\psi''|^q$  is  $(h, m)$ -convex modified of the second sense on  $[\rho, \frac{\sigma}{m}]$ , we have (11) and (12).

By substituting these values in (14), grouping the terms and proceeding as the previous Theorem, we obtain the inequality (13).

Then, this completes the proof.  $\square$

REMARK 11. By taking into account Remark 10, it is easy to obtain the Theorem 3.2 of [35] from the above result.

### 3. Conclusions

Although we have pointed out throughout the article that our results have special cases, some of which are found in the literature, we wanted to point out one more detail.

If we consider the function

$$w(\theta) = \begin{cases} \theta^{1+\alpha}, & \theta \in [0, \frac{1}{2}] \\ (1 - \theta)^{1+\alpha}, & \theta \in [\frac{1}{2}, 1], \end{cases}$$

from Lemma 1, we will obtain the equality

$$-\frac{1 + \alpha}{2^\alpha} \left[ \psi\left(\frac{3\rho + \sigma}{4}\right) + \psi\left(\frac{\rho + 3\sigma}{4}\right) \right] + \frac{2^{\alpha-1}\Gamma(\alpha + 2)}{(\sigma - \rho)^\alpha} \tag{15}$$

$$\times \left[ I_{\left(\frac{3\rho + \sigma}{4}\right)^+}^\alpha \psi\left(\frac{\rho + \sigma}{2}\right) + I_{\left(\frac{\rho + 3\sigma}{4}\right)^+}^\alpha \psi(\sigma) + I_{\left(\frac{3\rho + \sigma}{4}\right)^-}^\alpha \psi(\rho) + I_{\left(\frac{\rho + 3\sigma}{2}\right)^-}^\alpha \psi\left(\frac{\rho + \sigma}{2}\right) \right]$$

$$= \frac{(\sigma - \rho)^2}{8} \left\{ \int_0^{1/2} t^{1+\alpha} \left[ \psi'' \left( ta + (1-t) \frac{\rho + \sigma}{2} \right) + \psi'' \left( t \frac{\rho + \sigma}{2} + (1-t) \sigma \right) \right] dt \right. \\ \left. + \int_{1/2}^1 (1-\theta)^{1+\alpha} \left[ \psi'' \left( ta + (1-t) \frac{\rho + \sigma}{2} \right) + \psi'' \left( t \frac{\rho + \sigma}{2} + (1-t) \sigma \right) \right] dt \right\}.$$

This result is better than the one offered in Remark 7 and it is not reported in the literature.

In this way, we have the following inequality:

**THEOREM 5.** Let  $\psi : I = [\rho, \sigma] \rightarrow \mathbb{R}$  and  $\psi \in C^2(I^\circ)$  with  $\psi'' \in L_1(I)$ . If  $|\psi''|$  is a convex function ( $h(\theta) = \theta$  and  $s = m = 1$ ), then the inequality

$$\left| \frac{2^{\alpha-1} \Gamma(\alpha+2)}{(\sigma - \rho)^\alpha} \left[ I_{\left(\frac{3\rho+\sigma}{4}\right)^+}^\alpha \psi \left( \frac{\rho + \sigma}{2} \right) + I_{\left(\frac{\rho+3\sigma}{4}\right)^+}^\alpha \psi(\sigma) \right. \right. \\ \left. \left. + I_{\left(\frac{3\rho+\sigma}{4}\right)^-}^\alpha \psi(\rho) + I_{\left(\frac{\rho+3\sigma}{4}\right)^-}^\alpha \psi \left( \frac{\rho + \sigma}{2} \right) \right] - \frac{1+\alpha}{2^\alpha} \left[ \psi \left( \frac{3\rho + \sigma}{4} \right) + \psi \left( \frac{\rho + 3\sigma}{4} \right) \right] \right| \\ \leq \frac{(\sigma - \rho)^2}{2^{\alpha+5} (\alpha + 2)} \left[ \frac{(|\psi''(\rho)| - |\psi''(\sigma)|) (\alpha^2 + 6\alpha + 10)}{2(\alpha + 3)} \right. \\ \left. + (\alpha + 3) \left( \left| \psi'' \left( \frac{\rho + \sigma}{2} \right) \right| + |\psi''(\sigma)| \right) \right]. \quad (16)$$

holds  $\forall \alpha > 0$ .

*Proof.* From (15) and the module properties, we can write

$$\left| \frac{2^{\alpha-1} \Gamma(\alpha+2)}{(\sigma - \rho)^\alpha} \right. \\ \left. \times \left[ I_{\left(\frac{3\rho+\sigma}{4}\right)^+}^\alpha \psi \left( \frac{\rho + \sigma}{2} \right) + I_{\left(\frac{\rho+3\sigma}{4}\right)^+}^\alpha \psi(\sigma) + I_{\left(\frac{3\rho+\sigma}{4}\right)^-}^\alpha \psi(\rho) + I_{\left(\frac{\rho+3\sigma}{4}\right)^-}^\alpha \psi \left( \frac{\rho + \sigma}{2} \right) \right] \right. \\ \left. - \frac{1+\alpha}{2^\alpha} \left[ \psi \left( \frac{3\rho + \sigma}{4} \right) + \psi \left( \frac{\rho + 3\sigma}{4} \right) \right] \right| \\ \leq \frac{(\sigma - \rho)^2}{8} \left( \int_0^{1/2} t^{1+\alpha} \left| \psi'' \left( ta + (1-t) \frac{\rho + \sigma}{2} \right) + \psi'' \left( t \frac{\rho + \sigma}{2} + (1-t) \sigma \right) \right| dt \right. \\ \left. + \int_{1/2}^1 (1-\theta)^{1+\alpha} \left| \psi'' \left( ta + (1-t) \frac{\rho + \sigma}{2} \right) + \psi'' \left( t \frac{\rho + \sigma}{2} + (1-t) \sigma \right) \right| dt \right).$$

Since  $\psi$  is a convex for the integrals, we get:

$$|I_1| = \int_0^{1/2} t^{1+\alpha} \left| \psi'' \left( ta + (1-t) \frac{\rho + \sigma}{2} \right) + \psi'' \left( t \frac{\rho + \sigma}{2} + (1-t) \sigma \right) \right| dt \\ \leq (|\psi''(\rho)| - |\psi''(\sigma)|) \int_0^{1/2} \theta^{\alpha+2} d\theta + \left( \left| \psi'' \left( \frac{\rho + \sigma}{2} \right) \right| + |\psi''(\sigma)| \right) \int_0^{1/2} \theta^{\alpha+1} d\theta$$

$$= \frac{|\psi''(\rho)| - |\psi''(\sigma)|}{2^{\alpha+3}} + \frac{|\psi''\left(\frac{\rho+\sigma}{2}\right)| + |\psi''(\sigma)|}{2^{\alpha+2}},$$

and

$$\begin{aligned} |I_2| &= \int_{1/2}^1 (1-\theta)^{1+\alpha} \left| \psi''\left(t\rho + (1-t)\frac{\rho+\sigma}{2}\right) + \psi''\left(t\frac{\rho+\sigma}{2} + (1-t)\sigma\right) \right| dt \\ &\leq \int_{1/2}^1 (1-\theta)^{1+\alpha} \left[ \theta |\psi''(\rho)| + (1-\theta) \left| \psi''\left(\frac{\rho+\sigma}{2}\right) \right| \right. \\ &\quad \left. + \theta \left| \psi''\left(\frac{\rho+\sigma}{2}\right) \right| + (1-\theta) |\psi''(\sigma)| \right] d\theta \\ &= (|\psi''(\rho)| - |\psi''(\sigma)|) \int_{1/2}^1 (1-\theta)^{1+\alpha} \theta d\theta \\ &\quad + \left( \left| \psi''\left(\frac{\rho+\sigma}{2}\right) \right| + |\psi''(\sigma)| \right) \int_{1/2}^1 (1-\theta)^{1+\alpha} d\theta \\ &= \frac{(\alpha+4)(|\psi''(\rho)| - |\psi''(\sigma)|)}{2^{\alpha+3}(\alpha+2)(\alpha+3)} + \frac{|\psi''\left(\frac{\rho+\sigma}{2}\right)| + |\psi''(\sigma)|}{2^{\alpha+2}(\alpha+2)}. \end{aligned}$$

By summing the last inequality, we get:

$$\begin{aligned} &|I_1| + |I_2| \\ &\leq \frac{|\psi''(\rho)| - |\psi''(\sigma)|}{2^{\alpha+3}} + \frac{|\psi''\left(\frac{\rho+\sigma}{2}\right)| + |\psi''(\sigma)|}{2^{\alpha+2}} \\ &\quad + \frac{(\alpha+4)(|\psi''(\rho)| - |\psi''(\sigma)|)}{2^{\alpha+3}(\alpha+2)(\alpha+3)} + \frac{|\psi''\left(\frac{\rho+\sigma}{2}\right)| + |\psi''(\sigma)|}{2^{\alpha+2}(\alpha+2)} \\ &= \frac{|\psi''(\rho)| - |\psi''(\sigma)|}{2^{\alpha+3}} \left( 1 + \frac{\alpha+4}{(\alpha+2)(\alpha+3)} \right) + \frac{(|\psi''\left(\frac{\rho+\sigma}{2}\right)| + |\psi''(\sigma)|)(\alpha+3)}{2^{\alpha+2}(\alpha+2)}. \end{aligned}$$

Or, by multiplying both sides of the inequality by the expression  $\frac{(\sigma-\rho)^2}{8}$ , we get

$$\begin{aligned} &\frac{(\sigma-\rho)^2}{8} (|I_1| + |I_2|) \\ &\leq \frac{(\sigma-\rho)^2}{8} \left[ \frac{|\psi''(\rho)| - |\psi''(\sigma)|}{2^{\alpha+3}} \left( 1 + \frac{\alpha+4}{(\alpha+2)(\alpha+3)} \right) \right. \\ &\quad \left. + \frac{(|\psi''\left(\frac{\rho+\sigma}{2}\right)| + |\psi''(\sigma)|)(\alpha+3)}{(\alpha+2)2^{\alpha+2}} \right] \\ &= \frac{(\sigma-\rho)^2}{(\alpha+2)2^{\alpha+5}} \left[ \frac{(|\psi''(\rho)| - |\psi''(\sigma)|)(\alpha^2 + 6\alpha + 10)}{2(\alpha+3)} \right. \\ &\quad \left. + \left( \left| \psi''\left(\frac{\rho+\sigma}{2}\right) \right| + |\psi''(\sigma)| \right) (\alpha+3) \right] \end{aligned}$$

The proof is completed.  $\square$

Finally, we must remember that the presented results contain generalized inequalities which are valid for convex functions,  $m$ -convex functions,  $h$ -convex functions, and  $s$ -convex functions in the second sense, and defined in a closed interval of negative non-real numbers. It is clear that the problem of extending these results to the case of  $(h, m)$ -convex functions of the first type remains open.

## REFERENCES

- [1] M. A. ALI, J. E. NÁPOLES VALDÉS, A. KASHURI AND Z. ZHANG, *Fractional non conformable Hermite-Hadamard inequalities for generalized  $\phi$ -convex functions*, Fasciculi Mathematici, **64**, (2020), 5–16, <https://doi.org/10.21008/j.0044-4413.2020.0007>.
- [2] M. ALOMARI, M. DARUS AND S. S. DRAGOMIR, *New Inequalities Of Hermite-Hadamard Type for Functions Whose Second Derivatives absolute Values Are Quasi-Convex*, Tamkang Journal Of Mathematics, **41** 4, (2010) 353–359.
- [3] J. E. NÁPOLES VALDÉS AND B. BAYRAKTAR, *A note on Hermite-Hadamard integral inequality for  $(h, m)$ -convex modified functions in a generalized framework*, Submitted.
- [4] B. BAYRAKTAR, *Some new inequalities of Hermite-Hadamard type for differentiable Godunova-Levin functions via fractional integrals*, Konuralp Journal of Mathematics, **8** (1), (2020), 91–96.
- [5] B. BAYRAKTAR, *Some Integral Inequalities Of Hermite-Hadamard Type For Differentiable  $(s, m)$ -Convex Functions Via Fractional Integrals*, TWMS J. App. Eng. Math. **10** 3, (2020) 625–637.
- [6] B. BAYRAKTAR AND M. GÜRBÜZ, *On Some Integral Inequalities for  $(s, m)$ -Convex Functions*, TWMS Journal of Applied and Engineering Mathematics **10** 2 (2020), 288–295.
- [7] B. BAYRAKTAR, A. ATTAEV AND V. KUDAEV, *Some Generalized Hadamard Type Inequalities Via Fractional Integrals*, Russian Mathematics (Iz. VUZ), **65** (2), (2021), 1–14, <https://doi.org/10.3103/S1066369X21020018>.
- [8] S. BERMUDO, P. KÓRUS AND J. E. NÁPOLES VALDÉS, *On  $q$ -Hermite-Hadamard inequalities for general convex functions*, Acta Math. Hungar. **162** (2020), 364–374, <https://doi.org/10.1007/s10474-020-01025-6>.
- [9] W. W. BRECKNER, *Stetigkeitsaussagen für eine Klasse verallgemeinerter konvexer funktionen in topologischen linearen Räumen*, Publ. Inst. Math. **23** (1978), 13–20.
- [10] S. S. DRAGOMIR, R. P. AGARWAL, *Two inequalities for differentiable mappings and applications to special means of real numbers and trapezoidal formula*, Appl. Math. Lett., **11** (5), (1998) 91–95.
- [11] S. S. DRAGOMIR AND S. FITZPATRIK, *The Hadamard's inequality for  $s$ -convex functions in the second sense*, Demonstration Math., **32** (4) (1999), 687–696.
- [12] S. S. DRAGOMIR AND C. E. M. PEARCE, *Selected Topics on Hermite-Hadamard Inequalities*, RGMIA Monographs, Victoria University, (2000), available at [http://rgmia.vu.edu.au/monographs/hermite\\_hadamard.html](http://rgmia.vu.edu.au/monographs/hermite_hadamard.html).
- [13] S. S. DRAGOMIR, J. PECARIC AND L. E. PERSSON, *Some inequalities of Hadamard type*, Soochow J. Math., **21** (1995), 335–241.
- [14] J. E. HERNÁNDEZ HERNÁNDEZ, *On Some New Integral Inequalities Related With The Hermite-Hadamard Inequality via  $h$ -Convex Functions*, MAYFEB Journal of Mathematics, **4** (2017), 1–12.
- [15] H. HUDZIK AND L. MALIGRANDA, *Some remarks on  $s$ -convex functions*, Aequationes Math. **48** (1) (1994), 100–111.
- [16] S. HUSSAIN, F. AZHAR AND M. A. LATIF, *Generalized fractional Ostrowski type integral inequalities for logarithmically  $h$ -convex function*, The Journal of Analysis, **29** (2021), 1265–1278, <https://doi.org/10.1007/s41478-021-00310-z>.
- [17] P. M. GUZMÁN, J. E. NÁPOLES VALDÉS AND Y. GASIMOV, *Integral inequalities within the framework of generalized fractional integrals*, Fractional Differential Calculus, **11** (1) (2021), 69–84, <https://doi.org/10.7153/fdc-2021-11-05>.
- [18] I. IŞCAN, *A new generalization of some integral inequalities for  $(\alpha, m)$ -convex functions*, Math Sci **7** (2013) Article: 22, <https://doi.org/10.1186/2251-7456-7-22>.
- [19] W. J. LIU, Q. A. NGO AND V. N. HUY, *Several interesting integral inequalities*, J. Mathe. Ineq. **3** (2009), 201–212.

- [20] M. MATLOKA, *On Some Integral Inequalities For  $(h, m)$ -Convex Functions*, *Mathematical Economics*, **9** (16) (2013), 55–70.
- [21] V. G. MIHESAN, *A generalization of the convexity*, *Seminar on Functional Equations, Approx. and Convex.*, Cluj–Napoca (Romania) (1993).
- [22] H. MO AND X. SUI, *Hermite–Hadamard–type inequalities for generalized  $s$ -convex functions on real linear fractal set  $R^\alpha$ ,  $0 < \alpha < 1$* , *Math. Sci.* (2017) 11: 241–246, <https://doi.org/10.1007/s40096-017-0227-z>.
- [23] M. S. MOSLEHIAN, *Matrix Hermite–Hadamard type inequalities*, *Houston J. Math.*, **39** (1) (2013), 177–189.
- [24] M. MUDDASSAR, M. I. BHATTI AND W. IRSHAD, *Some new  $s$ -Hermite Hadamard type inequalities for differentiable functions and their applications*, *Proc. Pakistan Acad. Sci.* **49** (1) (2012) 9–17.
- [25] M. MUDDASSAR, M. I. BHATTI AND W. IRSHAD, *Generalisation of integral inequalities of Hermite–Hadamard type through convexity*, *Bull. Aust. Math. Soc.* **88** (2) (2014), 320–330.
- [26] J. E. NÁPOLES VALDÉS, F. RABOSI AND A. D. SAMANIEGO, *Convex Functions: Ariadne’s Thread Or Charlotte’s Spiderweb?*, *Advanced Mathematical Models & Applications* **5** (2) (2020), 176–191.
- [27] J. E. NÁPOLES VALDÉS, J. M. RODRÍGUEZ AND J. M. SIGARRETA, *On Hermite–Hadamard type inequalities for non–conformable integral operators*, *Symmetry*, **11**, (2019) 1108.
- [28] S. ÖZCAN, *Hermite–Hadamard type inequalities for  $m$ -convex and  $(\alpha, m)$ -convex functions*, *Journal of Inequalities and Applications* **2020**: 175 (2020), <https://doi.org/10.1186/s13660-020-02442-5>.
- [29] M. E. ÖZDEMİR, A. O. AKDEMİR AND E. SET, *On  $(h, m)$ -Convexity And Hadamard-Type Inequalities*, arXiv:1103.6163v1, 2011 – arxiv.org.
- [30] M. Z. SARIKAYA, A. SAGLAM AND H. YILDIRIN, *On Some Hadamard-Inequalities for  $h$ -convex Functions*, *Journal of Mathematical Inequalities*, **2** (3) (2008), 335–341.
- [31] M. Z. SARIKAYA, A. SAGLAM AND H. YILDIRIM, *New inequalities of Hermite–Hadamard type for functions whose second derivatives absolute values are convex and quasi-convex*, *International Journal of Open Problems in Computer Science and Mathematics* **5** (3) (2012), <https://doi.org/10.12816/0006114>.
- [32] G. TOADER, *Some generalizations of the convexity*, *Proceedings of the Colloquium on Approximation and Optimization*, University Cluj–Napoca, 329–338 (1985).
- [33] J. R. WANG, X. LI, M. FECKAN AND Y. ZHOU, *Hermite–Hadamard–type inequalities for Riemann–Liouville fractional integrals via two kinds of convexity*, *Applicable Analysis*, **92** (11) (2012), 1–13.
- [34] B. Y. XI AND F. QI, *Inequalities of Hermite–Hadamard type for extended  $s$ -convex functions and applications to means*, *J. Nonlinear Convex. Anal.* **16** (5) (2015), 873–890.
- [35] B. Y. XI, D. D. GAO AND F. QI, *Integral inequalities of Hermite–Hadamard type for  $(\alpha, s)$ -convex and  $(\alpha, s, m)$ -convex functions*, *Italian Journal Of Pure And Applied Mathematics*, **44**, (2020) 499–510.

(Received July 14, 2022)

Bahtiyar Bayraktar  
Bursa Uludağ University  
Department of Mathematics and Science Education  
Görükle Campus, 16059, Bursa, Turkey  
e-mail: bbayraktar@uludag.edu.tr

Juan E. Nápoles Valdés  
UTN-FRRE, French 414, Resistencia, Chaco 3500, Argentina  
e-mail: jnapoles@frre.utn.edu.ar

and

UNNE, FaCENA  
Ave. Libertad 5450, Corrientes 3400, Argentina  
e-mail: jnapoles@exa.unne.edu.ar