

MSC2020: 26A51, 26D10, 26D15

© *B. Bayraktar, J. E. Nápoles Valdés***NEW GENERALIZED INTEGRAL INEQUALITIES VIA (H, M) -CONVEX MODIFIED FUNCTIONS**

In this article, we establish several inequalities for (h, m) -convex maps, related to weighted integrals, used in previous works. Throughout the work, we show that our results generalize several of the integral inequalities known from the literature.

Keywords: Hermite–Hadamard inequality, Hölder inequality, power mean inequality, weighted integrals, (m, h) -convex functions

DOI: 10.35634/2226-3594-2022-60-01

Introduction

One of the basic concepts of a number of mathematical disciplines (function theory, optimization theory, theory of inequalities, etc.) is the concept of convexity. This concept is closely related to the estimation of the mean value of a function given on an interval. The definition of convexity of a function and the central inequality used to estimate the mean value of a function are given in the literature as follows:

Definition 0.1. The function $\phi: [v_1, v_2] \rightarrow \mathbb{R}$, is said to be convex, if we have

$$\phi(\varepsilon x + (1 - \varepsilon)y) \leq \varepsilon\phi(x) + (1 - \varepsilon)\phi(y)$$

$\forall x, y \in [v_1, v_2]$ and $\varepsilon \in [0, 1]$.

For convex functions on the interval $[v_1, v_2]$ at the end of the 19th century, the Hermite–Hadamard inequality

$$\phi\left(\frac{v_1 + v_2}{2}\right) \leq \frac{1}{v_2 - v_1} \int_{v_1}^{v_2} \phi(\varepsilon) d\varepsilon \leq \frac{\phi(v_1) + \phi(v_2)}{2}$$

widely known in the theory of inequalities, was established. Both inequalities hold in the reversed direction if ϕ is concave.

Today, in the literature there are quite a few different convexity classes of functions that are defined to refine, extend and improve estimates of the mean value of a function. A rather wide spectrum of convexity classes and their relations are given in [27].

Many researchers (see [2, 4–6, 11, 13, 14, 22, 25, 27, 28, 30, 33, 38–40] and references therein) have considered inequalities of the Hermite–Hadamard type with the use of integral operators for the purpose of generalization, refinement, and improvement.

For example, Bayraktar et al. in [4], Bayraktar and Özdemir in [5], by dividing the interval of integration into subintervals, obtained generalized inequalities for convex and s -convex functions were obtained in terms of fractional integration operators. Bessenyei and Páles in [6] studied generalized convex functions of high order, and, as a result, obtained extensions of the classical Hermite–Hadamard inequality. Using k -fractional integrals Butt et al. in [9] for the s -Godunov–Levin, quasi-convex and η -quasi-convex functions, new integral inequalities were obtained. Du et al. in [12] introduced the concept of generalized semi- (m, h) -preinvex functions

and obtained some estimates for the trapezoid inequality. Nàpoles et al. in [25] obtained new integral inequalities of the Hermite–Hadamard type for convex and quasi-convex functions in a generalized context. In [26], Nápoles and Bayraktar established these inequalities for h -convex functions in the framework of a certain generalized integral. Xi and Qi in [39] and Yildiz et al. in [40] obtained new integral inequalities for s -convex, (α, s) -convex and extended s -convex functions.

We presented the following definitions.

Definition 0.2. Let $h: [0, 1] \rightarrow \mathbb{R}$ be a nonnegative function, $h \neq 0$ and $\phi: I = [0, +\infty) \rightarrow [0, +\infty)$. If inequality

$$\phi(\varepsilon\xi + m(1 - \varepsilon)\varsigma) \leq h^s(\varepsilon)\phi(\xi) + m(1 - h^s(\varepsilon))\phi(\varsigma)$$

is fulfilled for all $\xi, \varsigma \in I$ and $\varepsilon \in [0, 1]$, where $m \in [0, 1]$, $s \in (0, 1]$, then it is said that function ϕ is (h, m) -convex modified of first type on I .

Definition 0.3. Let $h: [0, 1] \rightarrow \mathbb{R}$ nonnegative functions, $h \neq 0$ and $\phi: I = [0, +\infty) \rightarrow [0, +\infty)$. If inequality

$$\phi(\varepsilon\xi + m(1 - \varepsilon)\varsigma) \leq h^s(\varepsilon)\phi(\xi) + m(1 - h(\varepsilon))^s\phi(\varsigma)$$

is fulfilled for all $\xi, \varsigma \in I$ and $\varepsilon \in [0, 1]$, where $m \in [0, 1]$, $s \in [-1, 1]$, then it is said that function ϕ is (h, m) -convex modified of second type on I .

Remark 0.1. From Definitions 0.2 and 0.3 we can define $N_{h,m}^s[v_1, v_2]$, where $v_1, v_2 \in [0, +\infty)$, as the set of (h, m) -convex modified functions, for which $\phi(v_1) \geq 0$, characterized by the triple $(h(\varepsilon), m, s)$. Note that if:

1. $(h(\varepsilon), 0, 0)$, then we have increasing functions [8].
2. $(\varepsilon, 0, s)$, then we have s -starshaped functions [8].
3. $(\varepsilon, 0, 1)$, then we have starshaped functions [8].
4. $(\varepsilon, 1, 1)$, then ϕ is a convex function on $[0, +\infty)$ [8].
5. $(\varepsilon, m, 1)$, then ϕ is an m -convex function on $[0, +\infty)$ [36].
6. $(\varepsilon, 1, s)$ and $s \in (0, 1]$, then ϕ is an s -convex function on $[0, +\infty)$ [7, 15].
7. $(\varepsilon, 1, s)$ and $s \in [-1, 1]$, then ϕ is an extended s -convex function on $[0, +\infty)$ [39].
8. (ε, m, s) and $s \in (0, 1]$, then ϕ is an (s, m) -convex function on $[0, +\infty)$ [30].
9. $(\varepsilon^\alpha, m, 1)$ with $\alpha \in (0, 1]$, then ϕ is an (α, m) -convex function on $[0, +\infty)$ [23].
10. $(\varepsilon^\alpha, m, s)$ with $\alpha \in (0, 1]$, then ϕ is an s -(α, m)-convex function on $[0, +\infty)$ [38].
11. $(h(\varepsilon), m, 1)$, then we have a variant of an (h, m) -convex function on $[0, +\infty)$ [28].

Remark 0.2. In the different notions of convexity, if the direction of the inequality changes, it will be called concave.

In our work, we use the Euler Gamma functions Γ (see [32]) and Γ_κ (see [10]):

$$\begin{aligned}\Gamma(z) &= \int_0^\infty \varepsilon^{z-1} e^{-\varepsilon} d\varepsilon, \quad \operatorname{Re}(z) > 0, \\ \Gamma_\kappa(z) &= \int_0^\infty \varepsilon^{z-1} e^{-\varepsilon^\kappa/\kappa} d\varepsilon, \quad \kappa > 0.\end{aligned}$$

Here $\lim_{\kappa \rightarrow 1} \Gamma_\kappa(z) = \Gamma(z)$, $\Gamma_\kappa(z) = (\kappa)^{\frac{z}{\kappa}-1} \Gamma\left(\frac{z}{\kappa}\right)$ and $\Gamma_\kappa(z + \kappa) = z \Gamma_\kappa(z)$.

To encourage comprehension of the subject, we present the definition of Riemann–Liouville fractional integrals (with $0 \leq v_1 < \varepsilon < v_2 \leq \infty$). The first one is the classical Riemann–Liouville fractional integral.

Definition 0.4. Let $\phi \in L_1[v_1, v_2]$. Then the Riemann–Liouville fractional integrals of order $\alpha \in \mathbb{C}$, $\operatorname{Re}(\alpha) > 0$ are defined by (right and left respectively):

$$\begin{aligned} {}^\alpha I_{v_1^+} f(r) &= \frac{1}{\Gamma(\alpha)} \int_{v_1}^r (r - \varepsilon)^{\alpha-1} f(\varepsilon) d\varepsilon, \quad r > v_1, \\ {}^\alpha I_{v_2^-} f(r) &= \frac{1}{\Gamma(\alpha)} \int_r^{v_2} (\varepsilon - r)^{\alpha-1} f(\varepsilon) d\varepsilon, \quad r < v_2. \end{aligned}$$

Next we present the weighted integral operators, which will be the basis of our work.

Definition 0.5. Let $\phi \in L([v_1, v_2])$ and let w be a continuous and positive function, $w: [0, 1] \rightarrow [0, +\infty)$, with first order derivatives piecewise continuous on I . Then the weighted fractional integrals are defined by (right and left respectively):

$$\begin{aligned} J_{v_1^+}^w \phi(r) &= \int_{v_1}^r w' \left(\frac{\varepsilon - v_1}{r - v_1} \right) \phi(\varepsilon) d\varepsilon, \\ J_{v_2^-}^w \phi(r) &= \int_r^{v_2} w' \left(\frac{v_2 - \varepsilon}{v_2 - r} \right) \phi(\varepsilon) d\varepsilon, \end{aligned}$$

with $v_1 < r \leq v_2$.

Remark 0.3. To have a clearer idea of the amplitude of the Definition 0.5, let's consider some particular cases of the kernel w' :

1. Putting $w'(\varepsilon) \equiv 1$, we obtain the classical Riemann integral.
2. If $w'(\varepsilon) = \frac{\varepsilon(\alpha-1)}{\Gamma(\alpha)}$, then we obtain the right-sided Riemann–Liouville fractional integral, and the left-sided one can be obtained similarly.
3. With convenient kernel choices w' we can get the right and left k -Riemann–Liouville fractional integrals of [24], the right-sided fractional integrals of a function ϕ with respect to another function h on $[v_1, v_2]$ (see [2]), the right and left integral operator of [17], the right and left sided generalized fractional integral operators of [34] and the integral operators of [18] and [19], can also be obtained from above Definition by imposing similar conditions to w' .

Of course there are other known integral operators, fractional or not, that can be obtained as particular cases of the previous one, but we leave it to interested readers.

One of the most dynamic areas in Mathematical Sciences today is that of Integral Inequalities, mainly linked to the classical Hermite–Hadamard Inequality with the use of different integral, fractional or generalized operators. In particular, we can find different works with various particular operators (for example, see [1, 2, 9, 12, 16, 19, 21, 22, 26]). The advantage of using weighted integral operators is that it provides a unified and general formulation of various results that can be obtained as particular cases of ours. If we add to the above the definition of convexity used, we can realize the breadth and generality of the results obtained.

The main objective of this work is to obtain new variants of the classical Hermite–Hadamard Inequality, in the context of the weighted integrals of the Definition 0.5. We will show that our results complement or generalize several of those known from the literature.

§ 1. Results

To prove our theorems, we will need the following result.

L e m m a 1.1. *Let $\phi: [v_1, v_2] \rightarrow \mathbb{R}$ be a differentiable function, $v_1 < r \leq v_2$, $v_2 > 0$. If $\phi \in L^1([v_1, v_2])$, then we have:*

$$\begin{aligned} & \frac{(w(1) - w(0))(\phi(r) + \phi(v_1))}{2} - \frac{1}{2(r - v_1)} [J_{r-}^w \phi(v_1) + J_{v_1+}^w \phi(r)] \\ &= \frac{r - v_1}{2} \int_0^1 [w(1 - \varepsilon) - w(\varepsilon)] \phi'(\varepsilon v_1 + (1 - \varepsilon)r) d\varepsilon. \end{aligned}$$

P r o o f. From known properties we have for the right member:

$$\begin{aligned} & \int_0^1 [w(1 - \varepsilon) - w(\varepsilon)] \phi'(\varepsilon v_1 + (1 - \varepsilon)r) d\varepsilon \\ &= \int_0^1 w(1 - \varepsilon) \phi'(\varepsilon v_1 + (1 - \varepsilon)r) d\varepsilon - \int_0^1 w(\varepsilon) \phi'(\varepsilon v_1 + (1 - \varepsilon)r) d\varepsilon \\ &= I_1 - I_2. \end{aligned}$$

For the first integral we have:

$$\begin{aligned} I_1 &= \int_0^1 w(1 - \varepsilon) \phi'(\varepsilon v_1 + (1 - \varepsilon)r) d\varepsilon \\ & \left| \begin{aligned} u &= w(1 - \varepsilon) \implies du = -w'(1 - \varepsilon) d\varepsilon \\ dv &= \phi'(\varepsilon v_1 + (1 - \varepsilon)r) d\varepsilon \implies v = \frac{1}{v_1 - r} \phi(\varepsilon v_1 + (1 - \varepsilon)r) \end{aligned} \right| \\ &= \frac{w(1 - \varepsilon) \phi(\varepsilon v_1 + (1 - \varepsilon)r)}{v_1 - r} \Big|_0^1 + \frac{1}{v_1 - r} \int_0^1 w'(1 - \varepsilon) \phi(\varepsilon v_1 + (1 - \varepsilon)r) d\varepsilon \\ &= \frac{w(0) \phi(v_1) - w(1) \phi(r)}{v_1 - r} + \frac{1}{v_1 - r} \int_0^1 w'(1 - \varepsilon) \phi(\varepsilon v_1 + (1 - \varepsilon)r) d\varepsilon \end{aligned}$$

and changing the variable: $\varepsilon v_1 + (1 - \varepsilon)r = z$,

$$\begin{aligned} \varepsilon &= \frac{z - r}{v_1 - r} \implies d\varepsilon = \frac{dz}{v_1 - r}, \\ 1 - \varepsilon &= 1 - \frac{z - r}{v_1 - r} = \frac{v_1 - z}{v_1 - r} = \frac{z - v_1}{r - v_1}, \end{aligned}$$

we get

$$\begin{aligned} I_1 &= \frac{w(0) \phi(v_1) - w(1) \phi(r)}{v_1 - r} + \frac{1}{v_1 - r} \int_r^{v_1} w' \left(\frac{z - v_1}{r - v_1} \right) \phi(z) \frac{dz}{v_1 - r} \\ &= -\frac{w(0) \phi(v_1) - w(1) \phi(r)}{r - v_1} - \frac{1}{(v_1 - r)^2} \int_{v_1}^r w' \left(\frac{z - v_1}{r - v_1} \right) \phi(z) dz \\ &= \frac{w(1) \phi(r) - w(0) \phi(v_1)}{r - v_1} - \frac{1}{(r - v_1)^2} J_{v_1+}^w \phi(r). \end{aligned}$$

Similarly, for the I_2 , we have

$$\begin{aligned} I_2 &= \int_0^1 w(\varepsilon) \phi'(\varepsilon v_1 + (1 - \varepsilon)r) d\varepsilon \\ &= -\frac{w(1) \phi(v_1) - w(0) \phi(r)}{r - v_1} + \frac{1}{(r - v_1)^2} J_{r-}^w \phi(v_1). \end{aligned}$$

Thus, we get

$$\begin{aligned}
I_1 - I_2 &= -\frac{w(0)\phi(v_1) - w(1)\phi(r)}{r - v_1} - \frac{1}{(r - v_1)^2} J_{v_1+}^w \phi(r) \\
&\quad + \frac{w(1)\phi(v_1) - w(0)\phi(r)}{r - v_1} - \frac{1}{(r - v_1)^2} J_{r-}^w \phi(v_1) \\
&= \frac{[\phi(v_1) + \phi(r)] [w(1) - w(0)]}{r - v_1} - \frac{1}{(r - v_1)^2} [J_{v_1+}^w \phi(r) + J_{r-}^w \phi(v_1)].
\end{aligned}$$

From $I_1 - I_2$ we obtain the desired result. This completes the proof. $\square$

Remark 1.1. Putting $w'(\varepsilon) = \frac{1}{\Gamma(\alpha)} \varepsilon^{\alpha-1}$, we obtain the Lemma 3 of [29], if $r = v_2$ the above result reduces to the Lemma 2 of [35]. With the kernel $w'(\varepsilon) = \frac{1}{k\Gamma_k(\alpha)} \varepsilon^{\frac{\alpha}{k}-1}$ if we put $r = v_2$ we get Lemma 2.3 of [14] (also see Lemma 2.1 of [37]).

Remark 1.2. The main advantage of the previous result, over others known from the literature, in addition to the variety of “weights” $w'(\varepsilon)$ that we can consider (which can lead us to different integral operators), is the fact of the variability of the point $r \in (v_1, v_2)$, which obviously ensures a greater generality and strength of it.

Remark 1.3. If we consider, $r = \frac{v_1+v_2}{2}$ we obtain the following results, not reported in the literature.

Lemma 1.2. Let $\phi: [v_1, v_2] \rightarrow \mathbb{R}$ be a differentiable function, $v_1 < r \leq v_2$, $v_2 > 0$. If $\phi \in L^1([v_1, v_2])$, then we have:

$$\begin{aligned}
&\frac{2}{v_2 - v_1} \left\{ \left[w(0)\phi\left(\frac{v_1+v_2}{2}\right) - w(1)\phi(v_1) \right] - \left[w(1)\phi(v_1) - w(0)\phi\left(\frac{v_1+v_2}{2}\right) \right] \right\} \\
&- \frac{2}{(v_2 - v_1)^2} \left(J_{\frac{v_1+v_2}{2}-}^w \phi(v_1) + J_{v_1+}^w \phi\left(\frac{v_1+v_2}{2}\right) \right) \\
&= \int_0^1 [w(1-\varepsilon) - w(\varepsilon)] \phi' \left(\varepsilon v_1 + (1-\varepsilon) \frac{v_1+v_2}{2} \right) d\varepsilon.
\end{aligned}$$

Lemma 1.3. Let $\phi: [v_1, v_2] \rightarrow \mathbb{R}$ be a differentiable function, $v_1 < r \leq v_2$, $v_2 > 0$. If $\phi \in L^1([v_1, v_2])$, then we have:

$$\frac{\phi(v_1) + \phi\left(\frac{v_1+v_2}{2}\right)}{2} - \frac{1}{v_2 - v_1} \int_{v_1}^{\frac{v_1+v_2}{2}} \phi(\varepsilon) d\varepsilon = \frac{v_2 - v_1}{4} \int_0^1 (1 - 2\varepsilon) \phi\left(\varepsilon v_1 + \frac{v_1+v_2}{2}(1-\varepsilon)\right) d\varepsilon.$$

Considering $w(\varepsilon) = \varepsilon$, i. e., working with the classical Riemann integral.

Theorem 1.1. Let $0 \leq v_1 < v_2$, and $\phi \in C^1(v_1, v_2)$, where $v_1, v_2 \in \mathbb{R}$ such that $v_1 < r \leq v_2$, and ϕ be a function defined on the interval $[v_1, v_2]$. If $\phi' \in L[v_1, v_2]$ and $|\phi'|$ is a (h, m) -convex modified function of second type on $[v_1, v_2]$ for some fixed $m, s \in (0, 1]$, then the inequality

$$\begin{aligned}
&\left| A - \frac{1}{2(r - v_1)} (J_{v_1+}^w \phi(r) - J_{r-}^w \phi(v_1)) \right| \\
&\leq \frac{r - v_1}{2} \int_0^1 (w(1-\varepsilon) + w(\varepsilon)) \left[|\phi'(v_1)| h^s(\varepsilon) + m \left| \phi'\left(\frac{r}{m}\right) \right| (1 - h(\varepsilon))^s \right] d\varepsilon,
\end{aligned}$$

with $A = \frac{(w(1)-w(0))(\phi(r)+\phi(v_1))}{2}$, is fulfilled.

P r o o f. From Lemma 1.1 and taking into account the properties of modulus, we obtain

$$\begin{aligned}
& \left| v_1 - \frac{1}{2(r-v_1)} (J_{v_1+}^w \phi(r) - J_{r-}^w \phi(v_1)) \right| \\
& \leq \frac{r-v_1}{2} \left| \int_0^1 [w(1-\varepsilon) - w(\varepsilon)] \phi'(\varepsilon v_1 + (1-\varepsilon)r) d\varepsilon \right| \\
& = \frac{r-v_1}{2} \left\{ \int_0^1 w(1-\varepsilon) |\phi'(\varepsilon v_1 + (1-\varepsilon)r)| d\varepsilon + \int_0^1 w(\varepsilon) |\phi'(\varepsilon v_1 + (1-\varepsilon)r)| d\varepsilon \right\}
\end{aligned} \tag{1.1}$$

since $|\phi'|$ is an (h, m) -convex modified function of second type we have $|\phi'(\varepsilon v_1 + (1-\varepsilon)r)| \leq |\phi'(v_1)| h^s(\varepsilon) + m |\phi'(\frac{r}{m})| (1-h(\varepsilon))^s$, so

$$\begin{aligned}
& \left\{ \int_0^1 w(1-\varepsilon) |\phi'(\varepsilon v_1 + (1-\varepsilon)r)| d\varepsilon + \int_0^1 w(\varepsilon) |\phi'(\varepsilon v_1 + (1-\varepsilon)r)| d\varepsilon \right\} \\
& \leq |\phi'(v_1)| \int_0^1 w(1-\varepsilon) h^s(\varepsilon) d\varepsilon + m \left| \phi'(\frac{r}{m}) \right| \int_0^1 w(1-\varepsilon) (1-h(\varepsilon))^s d\varepsilon \\
& + |\phi'(v_1)| \int_0^1 w(\varepsilon) h^s(\varepsilon) d\varepsilon + m \left| \phi'(\frac{r}{m}) \right| \int_0^1 w(\varepsilon) (1-h(\varepsilon))^s d\varepsilon.
\end{aligned} \tag{1.2}$$

Substituting (1.2) into (1.1), rearranging and grouping, we get the desired inequality. $\square$

R e m a r k 1.4. If we take $|\phi'|$ s -convex, i. e., $h(u) = u$, $m = 1$ and $w(\varepsilon) = \varepsilon^\alpha$, we get Theorem 5 of [40].

T h e o r e m 1.2. Let $0 \leq v_1 < v_2$, and $\phi \in C^1(v_1, v_2)$, where $v_1, v_2 \in \mathbb{R}$ such that $v_1 < r \leq v_2$, and ϕ be a function defined on the interval $[v_1, v_2]$. If $\phi' \in L[v_1, v_2]$ and $|\phi'|^q$ is an (h, m) -convex modified function of second type on $[v_1, v_2]$ for some fixed $m, s \in (0, 1]$ and $q > 1$ with $\frac{1}{p} + \frac{1}{q} = 1$, then the inequality

$$\begin{aligned}
& \left| A - \frac{1}{2(r-v_1)} (J_{v_1+}^w \phi(r) - J_{r-}^w \phi(v_1)) \right| \\
& \leq \left(\frac{r-v_1}{2} \right) (B + C) \left[|\phi'(v_1)|^q \int_0^1 h^s(\varepsilon) d\varepsilon + m \left| \phi'(\frac{r}{m}) \right|^q \int_0^1 (1-h(\varepsilon))^s d\varepsilon \right]^{\frac{1}{q}}
\end{aligned}$$

holds with A as before and $B = \left(\int_0^1 w^p(1-\varepsilon) d\varepsilon \right)^{\frac{1}{p}}$ and $C = \left(\int_0^1 w^p(\varepsilon) d\varepsilon \right)^{\frac{1}{p}}$.

P r o o f. As in the proof of the previous result, we have

$$\begin{aligned}
& \left| A - \frac{1}{2(r-v_1)} (J_{v_1+}^w \phi(r) - J_{r-}^w \phi(v_1)) \right| \\
& \leq \frac{r-v_1}{2} \left| \int_0^1 [w(1-\varepsilon) - w(\varepsilon)] \phi'(\varepsilon v_1 + (1-\varepsilon)r) d\varepsilon \right| \\
& = \frac{r-v_1}{2} \left\{ \int_0^1 w(1-\varepsilon) |\phi'(\varepsilon v_1 + (1-\varepsilon)r)| d\varepsilon + \int_0^1 w(\varepsilon) |\phi'(\varepsilon v_1 + (1-\varepsilon)r)| d\varepsilon \right\}
\end{aligned} \tag{1.3}$$

Using the Hölder inequality on the two integrals of (1.3), we get $\left(\frac{1}{p} + \frac{1}{q} = 1 \right)$:

$$\begin{aligned}
& \int_0^1 w(1-\varepsilon) |\phi'(\varepsilon v_1 + (1-\varepsilon)r)| d\varepsilon \\
& \leq \left(\int_0^1 w^p(1-\varepsilon) d\varepsilon \right)^{\frac{1}{p}} \left(\int_0^1 |\phi'(\varepsilon v_1 + (1-\varepsilon)r)|^q d\varepsilon \right)^{\frac{1}{q}},
\end{aligned} \tag{1.4}$$

$$\begin{aligned}
& \int_0^1 w(\varepsilon) |\phi'(\varepsilon v_1 + (1 - \varepsilon)r)| d\varepsilon \\
& \leq \left(\int_0^1 w^p(\varepsilon) d\varepsilon \right)^{\frac{1}{p}} \left(\int_0^1 |\phi'(\varepsilon v_1 + (1 - \varepsilon)r)|^q d\varepsilon \right)^{\frac{1}{q}}, \tag{1.5}
\end{aligned}$$

using the (h, m) -convexity of $|\phi'|^q$ we obtain

$$\left(\int_0^1 |\phi'(\varepsilon v_1 + (1 - \varepsilon)r)|^q d\varepsilon \right)^{\frac{1}{q}} \leq \left(|\phi'(v_1)|^q h^s(\varepsilon) + m \left| \phi'\left(\frac{r}{m}\right) \right|^q (1 - h(\varepsilon))^s \right)^{\frac{1}{q}}, \tag{1.6}$$

so, using (1.6) in (1.4) and (1.5), and then in (1.3), rearranging, grouping and denoting $B = \left(\int_0^1 w^p(1 - \varepsilon) d\varepsilon \right)^{\frac{1}{p}}$ and $C = \left(\int_0^1 w^p(\varepsilon) d\varepsilon \right)^{\frac{1}{p}}$, we arrive at the desired inequality. $\square$

Remark 1.5. As in the previous Remark, this result yields Theorem 6 of [40]. If, in addition, we put $r = v_2$, then we obtain Corollary 1 of the above-mentioned work.

Remark 1.6. If we work with convex functions, i. e., $h(\varepsilon) = \varepsilon$, $s = m = 1$, $r = v_2$ and $w(\varepsilon) = \varepsilon$, then the above result becomes Theorem 2.3 from [11]. Theorem 1 of [20] is also a particular case of this result.

Theorem 1.3. Let $0 \leq v_1 < v_2$, and $\phi \in C^1(v_1, v_2)$, where $v_1, v_2 \in \mathbb{R}$ such that $v_1 < r \leq v_2$ and ϕ be a function defined on the interval $[v_1, v_2]$. If $\phi' \in L[v_1, v_2]$ and $|\phi'|^q$ is an (h, m) -convex modified function of second type on $[v_1, v_2]$ for some fixed $m, s \in (0, 1]$ and $q \geq 1$, then the inequality

$$\begin{aligned}
& \left| A - \frac{1}{2(r - v_1)} (J_{v_1+}^w \phi(r) - J_{r-}^w \phi(v_1)) \right| \\
& \leq \left(\frac{r - v_1}{2} \right) \cdot D \cdot \left(\int_0^1 w(1 - \varepsilon) \left(|\phi'(v_1)|^q h^s(\varepsilon) + m \left| \phi'\left(\frac{r}{m}\right) \right|^q (1 - h(\varepsilon))^s \right) d\varepsilon \right)^{\frac{1}{q}} \\
& + \left(\frac{r - v_1}{2} \right) \cdot E \cdot \left(\int_0^1 w(\varepsilon) \left(|\phi'(v_1)|^q h^s(\varepsilon) + m \left| \phi'\left(\frac{r}{m}\right) \right|^q (1 - h(\varepsilon))^s \right) d\varepsilon \right)^{\frac{1}{q}},
\end{aligned}$$

holds, with A as before and $D = \left(\int_0^1 w(1 - \varepsilon) d\varepsilon \right)^{1 - \frac{1}{p}}$ and $E = \left(\int_0^1 w(\varepsilon) d\varepsilon \right)^{1 - \frac{1}{p}}$.

Proof. As before

$$\begin{aligned}
& \left| v_1 - \frac{1}{2(r - v_1)} (J_{v_1+}^w \phi(r) - J_{r-}^w \phi(v_1)) \right| \\
& \leq \frac{r - v_1}{2} \left| \int_0^1 [w(1 - \varepsilon) - w(\varepsilon)] \phi'(\varepsilon v_1 + (1 - \varepsilon)r) d\varepsilon \right| \\
& = \frac{r - v_1}{2} \left\{ \int_0^1 w(1 - \varepsilon) |\phi'(\varepsilon v_1 + (1 - \varepsilon)r)| d\varepsilon + \int_0^1 w(\varepsilon) |\phi'(\varepsilon v_1 + (1 - \varepsilon)r)| d\varepsilon \right\}
\end{aligned}$$

Using now the well-known power mean inequality with modulus properties, we obtain:

$$\begin{aligned}
& \int_0^1 w(1 - \varepsilon) |\phi'(\varepsilon v_1 + (1 - \varepsilon)r)| d\varepsilon + \int_0^1 w(\varepsilon) |\phi'(\varepsilon v_1 + (1 - \varepsilon)r)| d\varepsilon \\
& \leq \left(\int_0^1 w(1 - \varepsilon) d\varepsilon \right)^{1 - \frac{1}{q}} \left(\int_0^1 w(1 - \varepsilon) |\phi'(\varepsilon v_1 + (1 - \varepsilon)r)|^q d\varepsilon \right)^{\frac{1}{q}} \\
& + \left(\int_0^1 w(\varepsilon) d\varepsilon \right)^{1 - \frac{1}{q}} \left(\int_0^1 w(\varepsilon) |\phi'(\varepsilon v_1 + (1 - \varepsilon)r)|^q d\varepsilon \right)^{\frac{1}{q}}.
\end{aligned}$$

From the (h, m) -convexity of $|\phi'|^q$, and a simple but tedious algebraic work, the proof of the Theorem is completed. $\square$

Remark 1.7. Assuming $|\phi'|^q$ s -convex and $w(\varepsilon) = \varepsilon^\alpha$, we get Theorem 7 of [40].

§ 2. Conclusions

In this paper, various extensions and generalizations of the classical Hermite–Hadamard Inequality have been established, in the context of recently defined weighted integral operators. Throughout it, various particular cases of the results obtained have been shown, which proves their breadth and strength. However, we do not want to conclude without pointing out two more aspects regarding the breadth of our results. Firstly, referring to the integral operator used, given that the weight function can include several known cases, we can add that if $w'(\varepsilon) = \varepsilon^{1-\frac{\alpha}{k}}$ (that is, we consider the k -integral of [15]), and $r = v_2$ the Lemma 1.1 reduces to Lemma 2.1 of [37], obviously many of the results of that work, can also be obtained from ours, considering convex functions. The second issue is the fact that the weighted operators used in our work can be used to derive new generalizations of other inequalities, for example, the Minkowski inequality (see, for example, [3]) where the weight $w'(\varepsilon) = \frac{1}{\Gamma(\alpha)}\varepsilon^{\alpha-1}$ is used. All of the above opens new and promising lines of work in this fruitful area.

The effectiveness of the method lies, fundamentally, in that we can provide a general working method that encompasses functions with first derivative, second or higher order derivatives, either with an integral operator, with two operators, on the same interval, on subintervals, without forgetting that it encompasses many of the known weights, used in various particular cases.

REFERENCES

1. Akdemir A. O., Butt S. I., Nadeem M., Ragusa M. A. New general variants of Chebyshev type inequalities via generalized fractional integral operators, *Mathematics*, 2021, vol. 9, issue 2, 122. <https://doi.org/10.3390/math9020122>
2. Akkurt A., Yildirim M. E., Yildirim H. On some integral inequalities for (k, h) -Riemann–Liouville fractional integral, *New Trends in Mathematical Science*, 2016, vol. 4, issue 2, pp. 138–146. <https://doi.org/10.20852/ntmsci.2016217824>
3. Aljaaidi T. A., Pachpatte D. New generalization of reverse Minkowski’s inequality for fractional integral, *Advances in the Theory of Nonlinear Analysis and its Applications*, 2021, vol. 5, no. 1, pp. 72–81. <https://doi.org/10.31197/atnaa.756605>
4. Bayraktar B., Attaev A. Kh., Kudaev V. Ch. Some generalized Hadamard type inequalities via fractional integrals, *Russian Mathematics*, 2021, vol. 65, issue 2, pp. 1–14. <https://doi.org/10.3103/S1066369X21020018>
5. Bayraktar B., Emin Özdemir M. Generalization of Hadamard-type trapezoid inequalities for fractional integral operators, *Ufa Mathematical Journal*, 2021, vol. 13, no. 1, pp. 119–130. <https://doi.org/10.13108/2021-13-1-119>
6. Bessenyei M., Páles Z. On generalized higher-order convexity and Hermite–Hadamard-type inequalities, *Acta Scientiarum Mathematicarum (Szeged)*, 2004, vol. 70, nos. 1–2, pp. 13–24.
7. Breckner W. W. Stetigkeitsaussagen für eine Klasse verallgemeinerter konvexer Funktionen in topologischen linearen Räumen, *Publications de l’Institut Mathématique. Nouvelle série*, 1978, vol. 23 (37), issue 43, pp. 13–20 (in German). <http://eudml.org/doc/257486>
8. Bruckner A. M., Ostrow E. Some function classes related to the class of convex functions, *Pacific Journal of Mathematics*, 1962, vol. 12, no. 4, pp. 1203–1215. <https://doi.org/10.2140/pjm.1962.12.1203>
9. Butt S. I., Bayraktar B., Umar M. Several new integral inequalities via k -Riemann–Liouville fractional integrals operators, *Problemy Analiza — Issues of Analysis*, 2021, vol. 10 (28), issue 1, pp. 3–22. <https://doi.org/10.15393/j3.art.2021.8770>
10. Díaz R., Pariguan E. On hypergeometric functions and Pochhammer k -symbol, *Divulgaciones Matemáticas*, 2007, vol. 15, no. 2, pp. 179–192.

11. Dragomir S. S., Agarwal R. P. Two inequalities for differentiable mappings and applications to special means of real numbers and to trapezoidal formula, *Applied Mathematics Letters*, 1998, vol. 11, issue 5, pp. 91–95. [https://doi.org/10.1016/S0893-9659\(98\)00086-X](https://doi.org/10.1016/S0893-9659(98)00086-X)
12. Du T., Awan M. U., Kashuri A., Zhao S. S. Some k -fractional extensions of the trapezium inequalities through generalized relative semi- (m, h) -preinvexity, *Applicable Analysis*, 2021, vol. 100, issue 3, pp. 642–662. <https://doi.org/10.1080/00036811.2019.1616083>
13. Farid G., Rehman A. U., Ain Q. U. k -fractional integral inequalities of Hadamard type for $(h - m)$ -convex functions, *Computational Methods for Differential Equations*, 2020, vol. 8, issue 1, pp. 119–140.
14. Farid G., Rehman A. U., Zahra M. On Hadamard-type inequalities for k -fractional integrals, *Nonlinear Functional Analysis and Applications*, 2016, vol. 21, no. 3, pp. 463–478. <http://nfaa.kyungnam.ac.kr/journal-nfaa/index.php/NFAA/article/view/902/832>
15. Hudzik H., Maligranda L. Some remarks on s -convex functions, *Aequationes Mathematicae*, 1994, vol. 48, issue 1, pp. 100–111. <https://doi.org/10.1007/BF01837981>
16. İlhan E. Analysis of the spread of Hookworm infection with Caputo–Fabrizio fractional derivative, *Turkish Journal of Science*, 2022, vol. 7, issue 1, pp. 43–52.
17. Jarad F., Abdeljawad T., Shah K. On the weighted fractional operators of a function with respect to another function, *Fractals*, 2020, vol. 28, no. 8, 2040011. <https://doi.org/10.1142/S0218348X20400113>
18. Jarad F., Uğurlu E., Abdeljawad T., Baleanu D. On a new class of fractional operators, *Advances in Difference Equations*, 2017, vol. 2017, issue 1, article 247. <https://doi.org/10.1186/s13662-017-1306-z>
19. Khan T. U., Khan M. A. Generalized conformable fractional operators, *Journal of Computational and Applied Mathematics*, 2019, vol. 346, pp. 378–389. <https://doi.org/10.1016/j.cam.2018.07.018>
20. Kırmacı U. S., Klaričić Bakula M., Özdemir M. E., Pečarić J. Hadamard-type inequalities for s -convex functions, *Applied Mathematics and Computation*, 2007, vol. 193, issue 1, pp. 26–35. <https://doi.org/10.1016/j.amc.2007.03.030>
21. Kızıl Ş., Ardiç M. A. Inequalities for strongly convex functions via Atangana–Baleanu integral operators, *Turkish Journal of Science*, 2021, vol. 6, issue 2, pp. 96–109.
22. Koca İ., Akçetin E., Yaprakdal P. Numerical approximation for the spread of SIQR model with Caputo fractional order derivative, *Turkish Journal of Science*, 2020, vol. 5, issue 2, pp. 124–139.
23. Miheşan V. G. A generalization of convexity, *Seminar on functional equations, approximation and convexity*, Cluj-Napoca, Romania: 1993.
24. Mubeen S., Habibullah G. M. k -fractional integrals and application, *International Journal of Contemporary Mathematical Sciences*, 2012, vol. 7, no. 2, pp. 89–94.
25. Nápoles Valdés J. E., Bayraktar B., Butt S. I. New integral inequalities of Hermite–Hadamard type in a generalized context, *Punjab University Journal Of Mathematics*, 2021, vol. 53, issue 11, pp. 765–777. <https://doi.org/10.52280/Pujm.2021.531101>
26. Nápoles Valdés J. E., Bayraktar B. On the generalized inequalities of the Hermite–Hadamard type, *Filomat*, 2021, vol. 35, issue 14, pp. 4917–4924. <https://doi.org/10.2298/FIL2114917N>
27. Nápoles Valdés J. E., Rabossi F., Samaniego A. D. Convex functions: Ariadne’s thread or Charlotte’s Spiderweb?, *Advanced Mathematical Models and Applications*, 2020, vol. 5, no. 2, pp. 176–191.
28. Özdemir M. E., Akdemir A. O., Set E. On $(h - m)$ -convexity and Hadamard-type inequalities, *Transylvanian Journal Of Mathematics And Mechanics*, 2016, vol. 8, no. 1, pp. 51–58.
29. Özdemir M. E., Dragomir S. S., Yildiz Ç. The Hadamard inequality for convex function via fractional integrals, *Acta Mathematica Scientia*, 2013, vol. 33B, issue 5, pp. 1293–1299. [https://doi.org/10.1016/S0252-9602\(13\)60081-8](https://doi.org/10.1016/S0252-9602(13)60081-8)
30. Park J. Generalization of Ostrowski-type inequalities for differentiable real (s, m) -convex mappings, *Far East Journal of Mathematical Sciences*, 2011, vol. 49, no. 2, pp. 157–171.
31. Pearce C. E. M., Pečarić J. Inequalities for differentiable mappings with application to special means and quadrature formulæ, *Applied Mathematics Letters*, 2000, vol. 13, issue 2, pp. 51–55. [https://doi.org/10.1016/S0893-9659\(99\)00164-0](https://doi.org/10.1016/S0893-9659(99)00164-0)
32. Rainville E. D. *Special functions*, New York: Macmillan, 1960.

33. Sahoo S.K., Ahmad H., Tariq M., Kodamasingh B., Aydi H., de la Sen M. Hermite–Hadamard type inequalities involving k -fractional operator for $(\bar{h}, m)$ -convex functions, *Symmetry*, 2021, vol. 13, issue 9, 1686. <https://doi.org/10.3390/sym13091686>
34. Sarıkaya M.Z., Ertuğral F. On the generalized Hermite–Hadamard inequalities, *Annals of the University of Craiova. Mathematics and Computer Science Series*, 2020, vol. 47, no. 1, pp. 193–213.
35. Sarıkaya M.Z., Set E., Yıldız H., Başak N. Hermite–Hadamard’s inequalities for fractional integrals and related fractional inequalities, *Mathematical and Computer Modelling*, 2013, vol. 57, issues 9–10, pp. 2403–2407. <https://doi.org/10.1016/j.mcm.2011.12.048>
36. Toader G. Some generalizations of the convexity, *Proceedings of the Colloquium on Approximation and Optimization*, Cluj-Napoca: University Cluj-Napoca, 1985, pp. 329–338.
37. Wu S., Iqbal S., Aamir M., Samraiz M., Younus A. On some Hermite–Hadamard inequalities involving k -fractional operators, *Journal of Inequalities and Applications*, 2021, vol. 2021, article number 32. <https://doi.org/10.1186/s13660-020-02527-1>
38. Xi B.-Y., Gao D.-D., Qi F. Integral inequalities of Hermite–Hadamard type for (α, s) -convex and (α, s, m) -convex functions, *Italian Journal Of Pure And Applied Mathematics*, 2020, no. 44, pp. 499–510.
39. Xi B.-Y., Qi F. Inequalities of Hermite–Hadamard type for extended s -convex functions and applications to means, *Journal of Nonlinear and Convex Analysis*, 2015, vol. 16, no. 5, pp. 873–890.
40. Yıldız Ç., Özdemir M. E., Önal H. K. Fractional integral inequalities via s -convex functions, *Turkish Journal of Analysis and Number Theory*, 2017, vol. 5, no. 1, pp. 18–22.

Received 06.04.2022

Accepted 29.09.2022

Bahtiyar Bayraktar, Candidate of Engineering, Assistant Professor, Faculty of Education, Bursa Uludağ University, Görükle Campus, 16059, Bursa, Turkey.

ORCID: <https://orcid.org/0000-0001-7594-8291>

E-mail: bbayraktar@uludag.edu.tr

Juan Eduardo Nápoles Valdés, PhD, Professor, Universidad Nacional del Nordeste, Ave. Libertad, 5450, Corrientes, 3400, Argentina;

Universidad Tecnológica Nacional, St. French, 414, Resistencia, Chaco, 3500, Argentina.

ORCID: <https://orcid.org/0000-0003-2470-1090>

E-mail: jnapoles@exa.unne.edu.ar

Citation: B. Bayraktar, J. E. Nápoles Valdés. New generalized integral inequalities via (h, m) -convex modified functions, *Izvestiya Instituta Matematiki i Informatiki Udmurtskogo Gosudarstvennogo Universiteta*, 2022, vol. 60, pp. 3–15.

Ключевые слова: неравенство Эрмита–Адамара, неравенство Гёльдера, степенное неравенство, взвешенные интегралы, (m, h) -выпуклые функции

УДК: 517.518, 517.218, 517.927

DOI: 10.35634/2226-3594-2022-60-01

В этой статье мы устанавливаем несколько неравенств для (h, m) -выпуклых отображений, связанных с взвешенными интегралами, которые использовались в предыдущих работах. На протяжении всей работы мы показываем, что наши результаты обобщают несколько известных из литературы интегральных неравенств.

СПИСОК ЛИТЕРАТУРЫ

1. Akdemir A. O., Butt S. I., Nadeem M., Ragusa M. A. New general variants of Chebyshev type inequalities via generalized fractional integral operators // *Mathematics*. 2021. Vol. 9. Issue 2. 122. <https://doi.org/10.3390/math9020122>
2. Akkurt A., Yildirim M. E., Yildirim H. On some integral inequalities for (k, h) -Riemann–Liouville fractional integral // *New Trends in Mathematical Science*. 2016. Vol. 4. Issue 2. P. 138–146. <https://doi.org/10.20852/ntmsci.2016217824>
3. Aljaaidi T. A., Pachpatte D. New generalization of reverse Minkowski’s inequality for fractional integral // *Advances in the Theory of Nonlinear Analysis and its Applications*. 2021. Vol. 5. No. 1. P. 72–81. <https://doi.org/10.31197/atnaa.756605>
4. Байрактаров Б. Р., Аттаев А. Х., Кудаяев В. Ч. Некоторые обобщенные неравенства типа Адамара через дробные интегралы // *Известия высших учебных заведений. Математика*. 2021. № 2. С. 3–18. <https://doi.org/10.26907/0021-3446-2021-2-3-18>
5. Bayraktar B., Emin Özdemir M. Generalization of Hadamard-type trapezoid inequalities for fractional integral operators // *Уфимский математический журнал*. 2021. Т. 13. Вып. 1. С. 119–130. <http://mi.mathnet.ru/ufa548>
6. Bessenyei M., Páles Z. On generalized higher-order convexity and Hermite–Hadamard-type inequalities // *Acta Scientiarum Mathematicarum (Szeged)*. 2004. Vol. 70. Nos. 1–2. P. 13–24.
7. Breckner W. W. Stetigkeitsaussagen für eine Klasse verallgemeinerter konvexer Funktionen in topologischen linearen Räumen // *Publications de l’Institut Mathématique. Nouvelle série*. 1978. Vol. 23 (37). Issue 43. P. 13–20 (in German). <http://eudml.org/doc/257486>
8. Bruckner A. M., Ostrow E. Some function classes related to the class of convex functions // *Pacific Journal of Mathematics*. 1962. Vol. 12. No. 4. P. 1203–1215. <https://doi.org/10.2140/pjm.1962.12.1203>
9. Butt S. I., Bayraktar B., Umar M. Several new integral inequalities via k -Riemann–Liouville fractional integrals operators // *Проблемы анализа — Issues of Analysis*. 2021. Т. 10 (28). Вып. 1. С. 3–22. <https://doi.org/10.15393/j3.art.2021.8770>
10. Díaz R., Pariguan E. On hypergeometric functions and Pochhammer k -symbol // *Divulgaciones Matemáticas*. 2007. Vol. 15. No. 2. P. 179–192.
11. Dragomir S. S., Agarwal R. P. Two inequalities for differentiable mappings and applications to special means of real numbers and to trapezoidal formula // *Applied Mathematics Letters*. 1998. Vol. 11. Issue 5. P. 91–95. [https://doi.org/10.1016/S0893-9659\(98\)00086-X](https://doi.org/10.1016/S0893-9659(98)00086-X)
12. Du T., Awan M. U., Kashuri A., Zhao S. S. Some k -fractional extensions of the trapezium inequalities through generalized relative semi- (m, h) -preinvexity // *Applicable Analysis*. 2021. Vol. 100. Issue 3. P. 642–662. <https://doi.org/10.1080/00036811.2019.1616083>

13. Farid G., Rehman A.U., Ain Q.U. k -fractional integral inequalities of Hadamard type for $(h - m)$ -convex functions // Computational Methods for Differential Equations. 2020. Vol. 8. Issue 1. P. 119–140.
14. Farid G., Rehman A.U., Zahra M. On Hadamard-type inequalities for k -fractional integrals // Nonlinear Functional Analysis and Applications. 2016. Vol. 21. No. 3. P. 463–478.
<http://nfaa.kyungnam.ac.kr/journal-nfaa/index.php/NFAA/article/view/902/832>
15. Hudzik H., Maligranda L. Some remarks on s -convex functions // Aequationes Mathematicae. 1994. Vol. 48. Issue 1. P. 100–111. <https://doi.org/10.1007/BF01837981>
16. İlhan E. Analysis of the spread of Hookworm infection with Caputo–Fabrizio fractional derivative // Turkish Journal of Science. 2022. Vol. 7. Issue 1. P. 43–52.
17. Jarad F., Abdeljawad T., Shah K. On the weighted fractional operators of a function with respect to another function // Fractals. 2020. Vol. 28. No. 8. 2040011.
<https://doi.org/10.1142/S0218348X20400113>
18. Jarad F., Uğurlu E., Abdeljawad T., Baleanu D. On a new class of fractional operators // Advances in Difference Equations. 2017. Vol. 2017. Issue 1. Article 247. <https://doi.org/10.1186/s13662-017-1306-z>
19. Khan T.U., Khan M.A. Generalized conformable fractional operators // Journal of Computational and Applied Mathematics. 2019. Vol. 346. P. 378–389. <https://doi.org/10.1016/j.cam.2018.07.018>
20. Kırmacı U.S., Klaričić Bakula M., Özdemir M.E., Pečarić J. Hadamard-type inequalities for s -convex functions // Applied Mathematics and Computation. 2007. Vol. 193. Issue 1. P. 26–35.
<https://doi.org/10.1016/j.amc.2007.03.030>
21. Kızıllı Ş., Ardiç M.A. Inequalities for strongly convex functions via Atangana–Baleanu integral operators // Turkish Journal of Science. 2021. Vol. 6. Issue 2. P. 96–109.
22. Koca İ., Akçetin E., Yaprakdal P. Numerical approximation for the spread of SIQR model with Caputo fractional order derivative // Turkish Journal of Science. 2020. Vol. 5. Issue 2. P. 124–139.
23. Miheşan V.G. A generalization of convexity // Seminar on functional equations, approximation and convexity. Cluj-Napoca, Romania: 1993.
24. Mubeen S., Habibullah G.M. k -fractional integrals and application // International Journal of Contemporary Mathematical Sciences. 2012. Vol. 7. No. 2. P. 89–94.
25. Nápoles Valdés J.E., Bayraktar B., Butt S.I. New integral inequalities of Hermite–Hadamard type in a generalized context // Punjab University Journal Of Mathematics. 2021. Vol. 53. Issue 11. P. 765–777.
<https://doi.org/10.52280/Pujm.2021.531101>
26. Nápoles Valdés J.E., Bayraktar B. On the generalized inequalities of the Hermite–Hadamard type // Filomat. 2021. Vol. 35. Issue 14. P. 4917–4924. <https://doi.org/10.2298/FIL2114917N>
27. Nápoles Valdés J.E., Rabossi F., Samaniego A.D. Convex functions: Ariadne’s thread or Charlotte’s Spiderweb? // Advanced Mathematical Models and Applications. 2020. Vol. 5. No. 2. P. 176–191.
28. Özdemir M.E., Akdemir A.O., Set E. On $(h - m)$ -convexity and Hadamard-type inequalities // Transylvanian Journal Of Mathematics And Mechanics. 2016. Vol. 8. No. 1. P. 51–58.
29. Özdemir M.E., Dragomir S.S., Yıldız Ç. The Hadamard inequality for convex function via fractional integrals // Acta Mathematica Scientia. 2013. Vol. 33B. Issue 5. P. 1293–1299.
[https://doi.org/10.1016/S0252-9602\(13\)60081-8](https://doi.org/10.1016/S0252-9602(13)60081-8)
30. Park J. Generalization of Ostrowski-type inequalities for differentiable real (s, m) -convex mappings // Far East Journal of Mathematical Sciences. 2011. Vol. 49. No. 2. P. 157–171.
31. Pearce C.E.M., Pečarić J. Inequalities for differentiable mappings with application to special means and quadrature formulæ // Applied Mathematics Letters. 2000. Vol. 13. Issue 2. P. 51–55.
[https://doi.org/10.1016/S0893-9659\(99\)00164-0](https://doi.org/10.1016/S0893-9659(99)00164-0)
32. Rainville E.D. Special functions. New York: Macmillan, 1960.
33. Sahoo S.K., Ahmad H., Tariq M., Kodamasingh B., Aydi H., de la Sen M. Hermite–Hadamard type inequalities involving k -fractional operator for $(\bar{h}, m)$ -convex functions // Symmetry. 2021. Vol. 13. Issue 9. 1686. <https://doi.org/10.3390/sym13091686>
34. Sarıkaya M.Z., Ertuğral F. On the generalized Hermite–Hadamard inequalities // Annals of the University of Craiova. Mathematics and Computer Science series. 2020. Vol. 47. No. 1. P. 193–213.
35. Sarıkaya M.Z., Set E., Yıldız H., Başak N. Hermite–Hadamard’s inequalities for fractional integrals and related fractional inequalities // Mathematical and Computer Modelling. 2013. Vol. 57. Issues 9–10.

P. 2403–2407. <https://doi.org/10.1016/j.mcm.2011.12.048>

36. Toader G. Some generalizations of the convexity // *Proceedings of the Colloquium on Approximation and Optimization*. Cluj-Napoca: University Cluj-Napoca, 1985. P. 329–338.
37. Wu S., Iqbal S., Aamir M., Samraiz M., Younus A. On some Hermite–Hadamard inequalities involving k -fractional operators // *Journal of Inequalities and Applications*. 2021. Vol. 2021. Article number 32. <https://doi.org/10.1186/s13660-020-02527-1>
38. Xi B.-Y., Gao D.-D., Qi F. Integral inequalities of Hermite–Hadamard type for (α, s) -convex and (α, s, m) -convex functions // *Italian Journal Of Pure And Applied Mathematics*. 2020. No. 44. P. 499–510.
39. Xi B.-Y., Qi F. Inequalities of Hermite–Hadamard type for extended s -convex functions and applications to means // *Journal of Nonlinear and Convex Analysis*. 2015. Vol. 16. No. 5. P. 873–890.
40. Yildiz Ç., Özdemir M.E., Önal H.K. Fractional integral inequalities via s -convex functions // *Turkish Journal of Analysis and Number Theory*. 2017. Vol. 5. No. 1. P. 18–22.

Поступила в редакцию 06.04.2022

Принята в печать 29.09.2022

Байрактаров Бахтияр Ризаевич, к. т. н., доцент, педагогический факультет, Университет Улудаг, 16059, Турция, Бурса, кампус Горукл.

ORCID: <https://orcid.org/0000-0001-7594-8291>

E-mail: bbayraktar@uludag.edu.tr

Наполес Вальдес Хуан Эдуардо, PhD, профессор, Северо-Восточный национальный университет, 3400, Аргентина, провинция Корриентес, г. Корриентес, пр. Свободы, 5450;

Национальный технологический университет, 3500, Аргентина, провинция Чако, г. Ресистенсия, ул. Французская, 414.

ORCID: <https://orcid.org/0000-0003-2470-1090>

E-mail: jnapoles@exa.unne.edu.ar

Цитирование: Б. Байрактар, Х. Э. Наполес Вальдес. Новые обобщенные интегральные неравенства через (h, m) -выпуклые модифицированные функции // *Известия Института математики и информатики Удмуртского государственного университета*. 2022. Т. 60. С. 3–15.