



How does a viviparous semifossorial lizard reproduce? *Ophiodes intermedius* (Squamata: Anguidae) from subtropical climate in the Wet Chaco region of Argentina

Martín A. Ortiz^a, Jorgelina M. Boretto^{b,*}, Nora R. Ibargüengoytía^b

^a Laboratorio de Herpetología, Facultad de Ciencias Exacta y Naturales y Agrimensura, Universidad Nacional del Nordeste, Av. Libertad 5460, 3400 Corrientes, Argentina

^b INIBIOMA (CONICET–Universidad Nacional del Comahue), Quintral 1250, 8400 San Carlos de Bariloche, Río Negro, Argentina

ARTICLE INFO

Article history:

Received 29 February 2016

Received in revised form 17 October 2016

Accepted 8 January 2017

Available online 16 January 2017

Keywords:

Asynchronous reproductive cycles

Gametogenesis

Fat body cycle

Glass snake

Sexual dimorphism

ABSTRACT

The best predictors of reproductive patterns are commonly associated with climate factors, but evolutionary history also plays an important role. The semifossorial and viviparous lizard *Ophiodes intermedius* from the Wet Chaco region of Argentina showed an annual cycle with asynchrony between males and females and an unusual pattern for subtropical climates, with vitellogenesis beginning in autumn, ovulation and copulation in spring, and births occurring in summer. Males exhibited annual variation of testicular size associated with spermatogenic activity, reaching their maximum gonadal activity in late summer (March), but sperm storage in the epididymis and/or deferent duct occurred throughout the year. Females showed an extended reproductive cycle beginning in mid-autumn (May) with vitellogenesis and finishing with births from late spring to mid-summer (December to February). Litter size varied from 4 to 9 offspring. Females reached sexual maturity at a larger snout–vent length and, overall, showed greater body size than males, while males exhibited larger heads than females. Fat body cycles indicated that females use lipid reserves to support vitellogenesis and embryo development, while males allocate lipid resources to the search for females, courtship and copulation rather than to gametogenesis. *Ophiodes intermedius* differed from other species of the genus in litter size, gestation period, timing of birth and the minimum size at sexual maturity, probably as a result of the influence of ecological, historical and phylogenetic factors.

© 2017 Elsevier GmbH. All rights reserved.

1. Introduction

A diversity of reproductive patterns have been registered among Squamates (Fitch, 1970, 1982; Vitt, 2015), ranging from continuous reproductive cycles (Vitt, 1983; Mojica et al., 2003; Watling et al., 2005; Lozano et al., 2015), characteristic of the tropics (Brown and Shine, 2006; Vitt, 2015), to seasonal reproductive patterns, present in a variety of habitats, from tropical to temperate areas (Fitch, 1970, 1982; Fitzgerald et al., 1999; Brown and Shine, 2006; Medina and Ibargüengoytía, 2010). Oviparous species frequently reproduce during spring and summer months (Fitch, 1982; Vitt, 2015), while for some viviparous lizards, gametogenesis, ovulation and mating occur in autumn and births during spring, representative of an autumn reproductive cycle (Guillette and Casas-Andreu,

1980, 1987; Guillette and Méndez-de la Cruz, 1993; Ramírez-Pinilla et al., 2009). There is a third pattern, the “partial autumn reproductive cycle”, in which vitellogenesis begins in autumn, ovulation and copulation occur in spring, and births take place in summer (Ramírez-Pinilla, 1992, 1995; Medina and Ibargüengoytía, 2010; Méndez-de la Cruz et al., 2013). The autumn reproductive patterns are frequently observed in species that inhabit high elevations in tropical and subtropical latitudes (Guillette and Casas-Andreu, 1980, 1987; Guillette and Méndez-de la Cruz, 1993; Ramírez-Pinilla, 1991, 1992, 1995; Ramírez-Bautista et al., 2014) or high latitudes at low elevations in Patagonia, Argentina (Boretto and Ibargüengoytía, 2009; Boretto et al., 2014).

The reproductive phenotypes are shaped by the interplay between the evolutionary history of a lineage (James and Shine, 1985; Dunham et al., 1988; Mesquita and Colli, 2010) and other biotic and abiotic factors (Vitt and Caldwell, 2014). For example, among abiotic factors, the most relevant are: temperature, photoperiod (Licht, 1971, 1973) and rainfall (Marion, 1982; Colli,

* Corresponding author.

E-mail address: borettojm@comahue-conicet.gob.ar (J.M. Boretto).

1991); and among biotic factors: food availability (Ballinger, 1977; Dunham, 1982) and the resulting energy storage as fat body masses (Hahn and Tinkle, 1965; Derickson, 1976; Bonnet et al., 1998; Warne et al., 2012), predation risk (Brown and Shine, 2006) and social structure (Ibargüengoytia and Cussac, 1999). However, the way these different factors affect fossorial lizards is still understudied (How and Shine, 1999; Resende and Nascimento, 2015).

Ophiodes Wagler, 1828 [RS1] is a Neotropical genus of the family Anguidae (Pyron et al., 2013) with only five recognized species (Cacciali and Scott, 2015), characterized by cryptozoic habits, a long, limbless cylindrical body, and a viviparous reproductive mode (Ceí, 1993; Pough et al., 2016). *Ophiodes intermedius* Boulenger, 1894 [RS2] is distributed in Bolivia, Paraguay, and the northern and central parts of Argentina (Ceí, 1993; Herrera et al., 2001), where it is present in grasslands (Ceí, 1993) and also in urban areas (Herrera et al., 2001). It feeds on insects and spiders (Cabrera, 2009), but little is known about its reproductive biology, probably due to its cryptozoic habits which impede any ecological study (Pizzatto, 2005). The investigation of the reproductive biology of *Ophiodes fragilis* (Pizzatto, 2005) and *Ophiodes* cf. *striatus* (Montechiaro et al., 2011) from the southeast of Brazil, and of *Ophiodes fragilis* (formerly *O. striatus*) from the Atlantic Forest of Brazil (Barros and Teixeira, 2007) revealed an annual reproductive cycle for females (Pizzatto, 2005; Montechiaro et al., 2011) and a continuous male cycle with sperm storage in the deferent duct until the mating season (Pizzatto, 2005; Montechiaro et al., 2011).

Here we present the male and female reproductive and fat body cycles, minimum adult size, litter size, and sexual dimorphism of *Ophiodes intermedius* from the Wet Chaco region of Argentina (subtropical climate). The reproductive biology of this semifossorial lizard is discussed in relation to environmental variables and compared with other Neotropical congeneric and sympatric species in order to better understand the possible evolutionary causes of the reproductive pattern observed.

2. Materials and methods

2.1. Study area and climate

O. intermedius was studied in the grasslands of Chaco and Corrientes provinces (Argentina), in the eastern district of the Chaco phytogeographic province known as Wet Chaco (Cabrera and Willink, 1973; Cabrera, 1976; Carnevali, 1994).

The climate in the northeast of Argentina is mainly warm and subtropical; there is no dry season although winter rainfall is significantly low (Cabrera, 1976; Bruniard, 1997). The average annual rainfall varies from 1000 to 1500 mm. The mean annual temperatures vary from 20 to 23 °C; the mean minimum temperatures occur in July and vary from 13 to 16 °C, while the mean maximum temperatures are recorded in January and vary from 26 to 28 °C (Cabrera, 1976; Carnevali, 1994; Bruniard, 1997). Frosts are uncommon and only registered in winter (Carnevali, 1994).

2.2. Sampling and laboratory methods

Males ($n = 16$) and females ($n = 25$) were collected from October 2009 until March 2014, with different capture techniques, such as visual search and capture by hand, pitfall traps with drift fences and artificial shelters. In addition, in order to complete our sample, we used 4 adult males (UNNEC: 0849, 6426, 6835, 7237), 5 adult females (UNNEC: 5031, 5714, 6794, 1185, 10056) and 1 juvenile (UNNEC: 5163) from the herpetological collection of the Universidad Nacional del Nordeste at Corrientes (UNNEC). The specimens were collected in the following localities: Rincon Chico Ombu, Ituzaingó Department (27°16'S, 56°25'W), Paso de

la Patria (27°19'20"S, 58°34'47"W) and Santa Ana de los Guácaras (27°21'S, 58°39'W), San Cosme Department, Mburucuyá, Mburucuyá Department (28°03'S, 58°14'W), Concepción, Concepción Department (28°24'S, 57°53'W), San Cayetano (27°33'22"S, 58°40'33'W) and Corrientes city (27°28'50"S, 58°44'24'W), Capital Department of Corrientes province, and Campo Largo, Independencia Department (27°26'49"S, 58°38'43'W) of Chaco province.

Lizards were sexed and then weighed (body mass, BM) to the nearest 0.01 g with a digital balance (Ohaus traveler scale TA320; Ohaus, Parsippany, NJ, USA) before euthanasia. Sexual dimorphism was studied considering the following variables measured with a digital vernier caliper to the nearest 0.01 mm: snout–vent length (SVL), head length (HL), head width (HW) and head height (HH) at the level of the interparietal scale, neck width (NW), hip width (HipW) measured as body width at the insertion of the vestigial hind legs, and maximum body width (BW). Specimens were euthanized by intraperitoneal administration of articaine and L-adrenaline, and the testes and ovaries were dissected out to be measured and processed. After fixation in Bouin's solution for 24 h, lizards were stored in 70% ethanol and deposited at the herpetological collection of the Universidad Nacional del Nordeste (UNNEC), Corrientes province (UNNEC: 10576, 10577, 13013–13051).

2.3. Male reproductive cycle

The male gonadal cycle was determined based on macro- and microscopic observations. Testes size (TS) was measured as the antero–posterior diameter using a digital caliper (± 0.1 mm). Gonads were dehydrated in a graded ethanol series and embedded in paraffin for 24 h in the oven at 52 °C. Following a histological conventional protocol, sections of 5 μ m were cut with a rotary microtome (Arcano KD-1508A; Zhejiang Jinhua Kedi Instrumental Equipment Co., Jinhua, China) and stained with hematoxylin and eosin.

Spermatogenic stages were determined based on the most advanced cell type present in the seminiferous tubules following Mayhew and Wright (1970), and cell types were recognised based on Uribe (2002). Five spermatogenic stages were defined: (I) only spermatogonia, (II) primary and secondary spermatocytes, (III) round spermatids and spermatids in the process of elongation, (IV) spermatozoa in the seminiferous tubules, and (V) regression with cellular debris and scarce spermatozoa in the tubular lumen (modified from Mayhew and Wright, 1970). The presence or absence of spermatozoa in the epididymis and/or ductus deferens was also registered. Minimum SVL at sexual maturity in males was determined based on the smallest specimen with spermatogenic activity (stages II–V) or spermatozoa in the epididymis and/or ductus deferens.

2.4. Female reproductive cycle

Female gonadal cycles were defined by the presence and number of vitellogenic follicles, oviductal embryos and corpora lutea. The sizes of follicles and/or embryos were defined as the diameter of the largest follicle and/or embryo for each female (Ibargüengoytia and Cussac, 1998) and measured with an ocular micrometer to the nearest 0.1 mm using an Olympus SZX9 stereoscopic microscope (Olympus, Tokyo, Japan). Pregnant females were classified (sensu Leyton et al., 1980) based on embryonic development as initial (from cleavage to somatic embryos), medium (curvate trunk, outlines of ears and eyes, and abundant yolk), and advanced (foetus with scales and pigmented skin). Litter size was determined by counting the number of ovulated oocytes or embryos in uterus (Ibargüengoytia and Cussac, 1998). Minimum SVL at sexual maturity for females was determined by considering the smallest female containing enlarged vitellogenic follicles, embryos in utero or cor-

pora lutea. Females without vitellogenic follicles and with oviducts with either no folds or only small folds were classified as juveniles. This method follows the definition of Ibagüengoytia and Cussac (1998): “Uterus type 1: folds, if present, are small and never spread over the entire uterus.”

2.5. Fat body cycles

Fat bodies were excised after the necropsy of each adult lizard, and weighed to the nearest 0.01 g using a digital balance (Ohaus traveler scale TA320).

2.6. Statistical analyses

Statistical analyses were conducted using Infostat (version 2011), SPSS (version 17.0), and Sigmaplot (version 10.0). Assumptions of normality and homogeneity of variance were tested with the Shapiro–Wilk test and with the Levene test, respectively. Dependence between variables was tested performing simple regression and Pearson correlation. When correlation between independent variables and SVL was found, residuals of the linear regression were used to perform further analysis (Ramírez-Bautista and Vitt, 1997).

To analyse the sexual dimorphism, we performed a Student's *t*-test to determine the difference between female and male SVL. Residuals of the regression of the ln-transformed morphological variables (BM, HL, HW, HH, HipW, and BW) and ln-transformed SVL were used in the stepwise discriminant analysis (based on the *p* value with $\alpha = 0.05$ and $\alpha = 0.10$ as input and output significance levels) to determine the variables that best explain the differences between the sexes. The significance level used was $P < 0.05$ for all statistical tests and results are presented as means $\pm$ standard deviation (SD).

3. Results

3.1. Body size, sexual maturity and sexual dimorphism

The minimum adult size for males was 94.5 mm SVL (corresponding to a specimen with spermatocytes in the seminiferous tubules=stage II). In females, the minimum adult size was 126.9 mm SVL (corresponding to an individual with vitellogenic follicles). SVL of adult males ranged from 94.5 to 177.0 mm (mean = 136.7 ± 22.2 , $n = 20$), and BM from 2.73 to 16.71 g (mean = 9.9 ± 4.36 g, $n = 14$), while SVL of adult females ranged from 126.9 to 205.8 mm (mean = 160.5 ± 18.7 , $n = 28$), and BM from 7.6 to 20.58 g (mean = 14.78 ± 3.77 g, $n = 24$). SVL of juveniles ranged from 55.6 to 118.9 mm (mean = 95.5 ± 22.2 , $n = 6$). Adult females were larger than adult males (SVL, Student's *t*-test; $T = -4.02$, $P < 0.001$, $n = 48$; Fig. 1), while males exhibited larger heads (HL, discriminant analysis; $\lambda = 0.684$, $X^2 = 9.695$, $df = 3$, $P < 0.002$, $n = 48$).

3.2. Male reproductive cycle

The relationship between male SVL and testicular size was positive (linear regression; $r^2 = 0.45$, $F_{1,18} = 14.78$, $P = 0.001$, $n = 20$). Therefore, adjusted testicular size increased at the end of spring (December), and reached the highest value in late summer (March), and for some males it remained high until mid-autumn (May; Fig. 2). Histological analyses showed that males exhibit different spermatogenic stages throughout the year. Males with primary and secondary spermatocytes (stage II, $n = 6$) were captured from late summer (March) to mid-spring (November), and males with round and elongated spermatid phases (stage III, $n = 7$) were captured from mid-winter (August) to late spring (December). Males with spermatozoa in the seminiferous tubules (stage IV, $n = 3$) were




Fig. 1. Box plot of the significant dimorphic traits ($P < 0.05$) of female and male *Ophiodes intermedius*.

found in late summer (March) and mid-autumn (May), while males with testicular regression (stage V, $n = 4$) were captured in spring (October, December) and in mid-summer (February; Fig. 3). 80% (16–20) of adult males with testes in different spermatogenic stages (stage II–V) exhibited spermatozoa in the epididymis and/or ductus deferens throughout the year.

3.3. Follicular size, litter size and female reproductive cycle

The vitellogenic cycle begins in mid-autumn (May), extending to early spring (October; Fig. 4). Pregnant females with early stage embryos (stage 1) were collected in early spring (September and October), with embryos at a later stage of development (stage 2) from early spring to early summer (October to January), and only one female with late stage embryos (stage 3) was captured (on December 13th). Additionally, one female with corpora lutea was collected in early summer (January), and adult non-reproductive females ($n = 11$) were captured throughout the year. Litter size ranged from 4 to 9 embryos (mean = 6.4 ± 1.44 , $n = 12$) and was not correlated with maternal SVL (Pearson correlation; $r = 0.32$, $P = 0.31$). The smallest lizard with a visible umbilical scar (neonate) measured 55.6 mm SVL and was captured in January.




Fig. 2. Variation in adjusted testicular size of adult males of *Ophiodes intermedius* throughout the year. Solid circles correspond to the residuals from snout–vent length versus testicular diameter regression.




Fig. 3. Spermatogenic stages of *Ophiodes intermedius* versus data. Absence (white circles) and presence (solid circles) of spermatozoa in the epididymis and/or deferent duct are indicated. The numbers in brackets are the numbers of observations.




Fig. 4. Female reproductive cycle of *Ophiodes intermedius*. The largest diameters of the ovarian follicles or embryos in adult females are represented: nonvitellogenic follicles (white circles), vitellogenic follicles (solid circles), nonvitellogenic follicles and corpora lutea (solid triangles), and embryos (asterisks). The numbers in brackets indicate the embryonic development stage as initial (1), medium (2), and advanced (3).

3.4. Fat body cycles

Male fat body mass showed a significant relationship with SVL (linear regression; $r^2 = 0.42$, $P = 0.006$, $n = 16$), but in females this relationship was not significant ($r^2 = 0.05$, $P = 0.312$, $n = 22$). There was also a positive and significant correlation between fat body mass and testicular diameter ($r^2 = 0.30$, $P = 0.028$, $n = 16$; Fig. 5A). In contrast, females exhibited a negative correlation between fat body mass and the sizes of follicles or embryos ($r^2 = -0.27$, $P = 0.014$, $n = 22$; Fig. 5B).

4. Discussion

O. intermedius from the subtropical climate of the Wet Chaco region shows an unusual reproductive pattern characterized by vitellogenesis starting in autumn and ending in spring. Male gametogenesis exhibits asynchrony with females but the storage of sperm in their reproductive tracts probably allows them to coordinate courtship and copulation in spring. The timing of reproductive events and the asynchrony between males and females observed in *O. intermedius* are common in viviparous lizards that live at high altitudes (Leyton et al., 1980; Guillelte and Casas-Andreu, 1980; Méndez-de la Cruz and Villagrán, 1998; Ramírez-Pinilla, 1991, 1992; Villagrán-Santa Cruz et al., 2009; Ramírez-Bautista et al., 2014) and latitudes (Jones et al., 1997; Boretto and Ibargüengoytia, 2009; Boretto et al., 2014).

The possible benefits of the autumn reproductive activity in *O. intermedius* reside in the early births (spring–summer) during the rainy season, when temperature and food availability are high. Early births would allow newborns to grow and store energy before the next winter, increasing their survival probability as has been suggested for other viviparous species living in harsh environments (Guillelte and Bearce, 1986; Boretto and Ibargüengoytia, 2006, 2009; Ramírez-Pinilla et al., 2009; Boretto et al., 2007, 2014; Cabezas Cartes et al., 2010). In addition, in viviparous species, embryonic development during spring and summer months can be linked to the need for suitable environmental temperatures for the normal development of embryos (Almeida-Santos and Salomão, 2002), since temperature stability during development is considered a factor in preventing abnormalities and shortening the development period (Hubert, 1985).

The asynchronous gametogenic activity between males and females, as observed in *O. intermedius*, has been attributed to the different timing needs for the maturation of male and female gonads, the different energy requirements for gametogenesis of each sex, and the need to complete the reproductive cycle in short activity seasons (Jones et al., 1997; Boretto and Ibargüengoytia, 2006; Boretto et al., 2007, 2015; Cabezas Cartes et al., 2010; Mouton et al., 2012). In asynchronous cycles, sperm storage is obligatory (Mouton et al., 2012), either in the reproductive tract of females (Ramírez-Bautista et al., 1998; Sever and Hopkins, 2004; Ortega-León et al., 2009) or of males (Guillelte and Casas-Andreu, 1980; Jones et al., 1997; Boretto et al., 2007; Cabezas Cartes et al., 2010).

The presence of a reproductive pattern characteristic of cold temperate environments in *O. intermedius* that lives in a subtropical climate in the Wet Chaco region could be a result of the retention of an ancestral reproductive pattern evolved in a cold climate, as has been reported for a diversity of snake species from tropical and subtropical North America (Quinn, 1979; Jackson and Franz, 1981; Johnson et al., 1982) and South America (Almeida-Santos and Salomão, 2002; Almeida-Santos et al., 2006; Marques et al., 2013; Barros et al., 2012). In addition, even though *O. intermedius* lives in a warmer environment than its ancestors, the male gonadal cycle and the temporally dissociated mating may be a result of their semifos-

social habit if moist and cool conditions prevail in their microhabitat during the coldest months of the year.

According to molecular data, the genus *Ophiodes* may have originated about 21–22 million years ago (mya), in the early Miocene (Wiens and Slingluff, 2001). The evidence suggests that orogenic and climatic conditions were very unstable around the Oligocene–Miocene boundary (approximately 23 mya). The concentration of oxygen isotopes recorded in the early Miocene makes evident a short period of glaciation after which the climate became warmer until the mid-Miocene (17–15 mya), when a gradual drop of global temperatures began (Miller et al., 1991; Zachos et al., 2001). In South America, the cooling and aridification processes from the mid to late Miocene were associated with global climatic and local orogenic events (uplifting of the Andes), and a marine transgression that covered vast plains of South America (Janis, 1993; Webb, 1995; Ortiz-Jaureguizar and Cladera, 2006; Rossetti et al., 2013). These events led to the expansion of open habitats (grasslands and savannas) at the expense of woodlands (Retallack, 2001; Strömberg, 2011). In this relatively cool and dry scenario that may have prevailed in the past, the species of the genus *Ophiodes* may have developed their viviparous reproductive mode and their present reproductive pattern. The origin and evolution of viviparity in Squamata have been commonly associated with cold environments of high latitudes or altitudes, gradually developed from an oviparous ancestor by the progressive increase of the retention period of the egg, shell reduction, and development of some kind of primitive placenta for the exchange of water and gases (Tinkle and Gibbons, 1977; Blackburn, 1982, 2005, 2015; Shine, 1985; Guillelte, 1993). Pyron and Burbrink (2014) suggested that viviparity could be an ancestral condition in squamate reptiles; however, more recent reviews mention a variety of morphological, physiological and ecological factors that refute this hypothesis (Griffith et al., 2015), and the reversal from viviparity to oviparity could only have occurred in a few squamate species (King and Lee, 2015). *O. intermedius* females of the Wet Chaco produce oocytes with a large amount of yolk, resulting, after fertilization, in macrolecithal eggs. The embryos are surrounded by a thin transparent membrane sac, without the calcareous deposits of oviparous species. The abundant deposition of yolk in the oocytes suggests that embryo nutrition depends mainly on egg yolk, which is consistent with other species of this genus (Pizzatto, 2005; Barros and Teixeira, 2007). However, we cannot discard the possibility that there is also an exchange of organic and inorganic substances as has been found in other lizards (Stewart and Castillo, 1984; Stewart and Thompson, 1993, 2000; Crocco et al., 2008).

Since ovulation in *O. intermedius* occurs in September, one would expect to find storage of sperm not only in males, but also in females, given the fact that we recorded bite marks with a “V” shape, typically produced during mating (Anderson and Vitt, 1990), on the head of *O. intermedius* females captured in May, July and October. These marks indicate that mating occurs during the months of vitellogenic activity (May to November), when males show spermatocytes and spermatids in testes. It has been suggested that storage of sperm in males may indicate a prolonged period of mating, and may be obligatory if the optimal time for spermatogenesis and oogenesis occurs at different times of the year (Mitchell and Zug, 1984; Birkhead and Møller, 1993; Almeida-Santos et al., 2006), while sperm storage in females could promote sperm competition, causing them to be inseminated by several males (Birkhead and Møller, 1993; Adams et al., 2005; Barros et al., 2012). The storage of sperm in the ductus deferens seems to be a character shared by males of the genus *Ophiodes*; however, in *O. intermedius* from the Wet Chaco, the spermatogenic activity is seasonal, while the male cycle in *O. fragilis* and *O. cf. s. triatus* from subtropical regions of the Atlantic Forest in Brazil is continuous (Pizzatto, 2005; Montechiaro et al., 2011).




Fig. 5. Fat body annual cycle of *Ophiodes intermedius* adults. (A) Monthly means of the residuals obtained from the linear regressions between fat body mass and snout–vent length (triangles), and between means of testicular diameter and snout–vent length (circles). (B) Monthly means of fat bodies (triangles), and means of the largest follicles or embryos (circles) are also indicated. Bars indicate the standard error.

Reproductive asynchrony between males and females might favour the development of dimorphic traits that increase mating ability (Ramírez-Bautista et al., 2002). *O. intermedius* males exhibited a head size proportionately larger than females, which could be related to intrasexual competition. This could mean an advantage over other males during agonistic encounters for the access to females (Vitt and Cooper, 1985; Anderson and Vitt, 1990; Andersson, 1994), and might also relate to the advantage of better apprehension of the female during copulation (Anderson and Vitt, 1990; Hews, 1990). Probably, this is the case in *O. intermedius*, as evidenced by the presence of bite marks of copulation in females. Yet, larger heads in males have also been linked to niche partition between the sexes (Schoener, 1967; Schoener and Gorman, 1968; Presst, 1994; Herrel et al., 1999). Meanwhile, females showed a greater SVL, as has been observed in other *Ophiodes* species (Pizzatto, 2005; Barros and Teixeira, 2007; Montechiaro et al., 2011), probably as a result of fecundity selection, as has been reported in other species (Shine, 1988). However, while in other *Ophiodes* species the litter size depends on female SVL (*O. fragilis*, Santa Teresa population, Barros and Teixeira, 2007; *O. cf. striatus*, Montechiaro et al., 2011), litter size was independent of female body size in *O. intermedius* (present study) and *O. fragilis* from the

southeast of Brazil (Pizzatto, 2005). Nonetheless, other factors can interfere with the relationship between litter size and maternal body size. For example, the nutritional conditions and the amount of lipid storage in females can also influence litter and/or offspring size (Doughty and Shine, 1998; Bonnet et al., 2001).

Our study of fat body mass in *O. intermedius* revealed a pattern of energy allocation linked to reproduction, as has been described in a diversity of lizard species (Rocha, 1992; Guillette and Bearce, 1986; Ramírez-Bautista et al., 2009). The increment of fat body masses with spermatogenesis suggests that males probably allocate more resources to the search for females, courtship and copulation during the mating period than to the spermatogenic activity (energetically less expensive), while females probably use lipid reserves to support vitellogenesis and embryonic development, as previously documented for other lizards (Avery, 1970; Goldberg, 1972; Guillette and Casas-Andreu, 1981; Guillette and Bearce, 1986; Ramírez-Bautista et al., 2009).

Female cycles and reproductive parameters of the species of the genus *Ophiodes* show variations in the minimum size at sexual maturity, the vitellogenic cycle, the timing of ovulation and births, and the embryonic development period (Table 1). In particular, as latitude increases, the timing of births in *Ophiodes* species

Table 1
Overview of reproductive characteristics of the investigated *Ophiodes* species. Comparisons among reproductive season, female size at sexual maturity, mean adult female size, litter size, and birthing period in different biomes and localities. Mean values are presented with ± 1 SD, and with ranges in parentheses.

Species	Biome	Reproductive period	Female SVL (mm) at sexual maturity	Female mean size (SVL, mm)	Litter size	Birthing period	Locality	References
<i>Ophiodes intermedius</i>	Wet Chaco	May–January	126.9	157.0 $\pm$ 21.56 (112.7–205.8)	6.4 $\pm$ 1.44 (4–9)	December–January	Chaco and Corrientes, northeastern Argentina	Present study
<i>Ophiodes fragilis</i>	Atlantic Forest	February–November	156.0	192.5 $\pm$ 28 (56–300)	7.5 $\pm$ 2.3 (5–13)	August–November	São Paulo and Paraná, southeastern Brazil	Pizzatto (2005)
<i>Ophiodes fragilis</i>	Atlantic Forest	March–November	161.0	179.7 $\pm$ 25.5 (111.0–235.0)	6.0 $\pm$ 2.0 (3–11)	November	Espirito Santo, southeastern Brazil	Barros and Teixeira (2007)
<i>Ophiodes cf. striatus</i>	Atlantic Forest	February–December	107.0	161.0 $\pm$ 25.7 (107–220)	7.9 $\pm$ 3.8 (2–14)	December	Rio Grande do Sul, Santa Catarina and Paraná, southeastern Brazil	Montechiaro et al. (2011)
<i>Ophiodes striatus</i>	Pampa	–	145	218.0 $\pm$ 38.0 (145–252)	10.5 $\pm$ 7.6 (5–24)	December	Rio Grande do Sul, southeastern Brazil	Leitão (1973)

shifts from early spring to early or mid-summer (Table 1), probably related to temperature and rainfall patterns (Montechiaro et al., 2011). *O. intermedius* deviates from the usual reproductive pattern of this lineage, showing an asynchrony between male and female reproductive cycles, which is an unusual pattern in subtropical climates. This reproductive style differs from that of sympatric lizard species previously studied, like *Tropidurus catalanensis* (formerly *T. torquatus*, Ortiz et al., 2014) and *Kentropyx viridistriga* (Ortiz et al., 2016). The comparative analysis showed that *O. intermedius* from the Wet Chaco exhibits reproductive characteristics and sexual dimorphism that reveal the influence of ecological, historical and phylogenetic factors.

Acknowledgements

We thank J. L. Acosta, E. Etchepare, J. Valdés, R. Aguirre, D. Aguiar, S. Palomas, A. C. Falcione and M. R. Ingaramo for their assistance during field work. We are grateful to the anonymous reviewers for their critical reviews and their insightful comments on the manuscript. This study was supported by a research grant from the Secretaría General de Ciencia y Técnica (SGCyT) and the Consejo Nacional de Investigaciones Científicas y Técnicas (PIP11220120100676; RD0484-13). We also thank the Dirección de Fauna of Corrientes for the collecting permits.

References

- Adams, E.M., Jones, A.G., Arnold, S.J., 2005. Multiple paternity in a natural population of a salamander with long-term sperm storage. *Mol. Ecol.* 14, 1803–1810.
- Almeida-Santos, S.M., Salomão, M.G., 2002. Reproduction in Neotropical pitvipers, with an emphasis on species of the genus *Bothrops*. In: Schuett, G.W., Höggren, M., Douglas, M.E., Greene, H.W. (Eds.), *Biology of the Vipers*. Eagle Mountain Publishing, Eagle Mountain, UT, pp. 445–462.
- Almeida-Santos, S.M., Pizzatto, L., Marques, O.A.V., 2006. Intra-sex synchrony and inter-sex coordination in the reproductive timing of the Atlantic coral snake *Micrurus corallines* (Elapidae) in Brazil. *Herpetol. J.* 16, 371–376.
- Anderson, R.A., Vitt, L.J., 1990. Sexual selection versus alternative causes of sexual dimorphism in teiid lizards. *Oecologia* 84, 145–157.
- Andersson, M.B., 1994. *Sexual Selection*. Princeton University Press, Princeton.
- Avery, R.A., 1970. Utilization of caudal fat by hibernating common lizards, *Lacerta vivipara*. *Comp. Biochem. Physiol.* 37, 119–121.
- Ballinger, R.E., 1977. Reproductive strategies: food availability as a source of proximal variation in a lizard. *Ecology* 58, 628–635.
- Barros, E.H., Teixeira, R.L., 2007. Diet and fecundity of the glass-lizard, *Ophiodes striatus* (Sauria, Anguillidae) from the Atlantic Forest in southeastern Brazil. *Bol. Mus. Biol. Mello Leitão (N. Sér.)* 22, 11–23.
- Barros, V.A., Sueiro, L.R., Almeida-Santos, S.M., 2012. Reproductive biology of the Neotropical rattlesnake *Crotalus durissus* from northeastern Brazil: a test of phylogenetic conservatism of reproductive patterns. *Herpetol. J.* 22, 97–104.
- Birkhead, T.R., Möller, A.P., 1993. Sexual selection and the temporal separation of reproductive events: sperm storage data from reptiles, birds and mammals. *Biol. J. Linn. Soc.* 50, 295–311.
- Blackburn, D.G., 1982. Evolutionary origins of viviparity in the Reptilia. I. Sauria. *Amphibia-Reptilia* 3, 185–205.
- Blackburn, D.G., 2005. Amniote perspectives on the evolutionary origins of viviparity and placentation. In: Grier, H.J., Uribe, M.C. (Eds.), *Viviparous Fishes*. New Life Publications, Homestead, FL, pp. 301–322.
- Blackburn, D.G., 2015. Evolution of vertebrate viviparity and specializations for fetal nutrition: a quantitative and qualitative analysis. *J. Morphol.* 276, 961–990.
- Bonnet, X., Bradshaw, D., Shine, R., 1998. Capital versus income breeding: an ectothermic perspective. *Oikos* 83, 333–342.
- Bonnet, X., Naulleau, G., Shine, R., Lourdaux, O., 2001. Short-term versus long-term effects of food intake on reproductive output in a viviparous snake, *Vipera aspis*. *Oikos* 92, 297–308.
- Boretto, J.M., Ibagüengoytia, N.R., 2006. Asynchronous spermatogenesis and biennial female cycle of the viviparous lizard *Phymaturus antofagastensis* (Liolaemidae): reproductive responses to high altitudinal and temperature climate of Catamarca, Argentina. *Amphibia-Reptilia* 27, 25–36.
- Boretto, J.M., Ibagüengoytia, N.R., 2009. *Phymaturus* of Patagonia, Argentina: reproductive biology of *Phymaturus zapalensis* (Liolaemidae) and a comparison of sexual dimorphism within the genus. *J. Herpetol.* 43, 93–104.
- Boretto, J.M., Ibagüengoytia, N.R., Acosta, J.C., Blanco, G.M., Villavicencio, J., Marín, J.A., 2007. Reproductive biology and sexual dimorphism of a high-altitude population of the viviparous lizard *Phymaturus punae* from the Andes in Argentina. *Amphibia-Reptilia* 28, 427–432.
- Boretto, J.M., Cabezas Cartes, F., Tappari, F., Méndez-de la Cruz, F., Sinervo, B., Scolaro, A.J., Ibagüengoytia, N.R., 2014. Reproductive biology of *Phymaturus spectabilis*

- (Liolaemidae): females skip reproduction in cold and harsh environments of Patagonia, Argentina. *Herpetol. Conserv. Biol.* 9, 170–180.
- Boretto, J.M., Cabezas Cartes, F., Ibagüengoytia, N.R., 2015. Energy allocation to growth and reproduction in a viviparous lizard endemic to the highlands of the Andes, Argentina. *J. Zool.* 297, 77–86.
- Brown, G.P., Shine, R., 2006. Why do most tropical animals reproduce seasonally? Testing hypotheses on an Australian snake. *Ecology* 87, 133–143.
- Bruniari, E., 1997. Atlas Geográfico de la Provincia de Corrientes. Tomo I: El Medio Natural. Geografía 8. Facultad de Humanidades. Univ. Nac. Nordeste, Resistencia, Chaco, Argentina.
- Cabezas Cartes, F., Boretto, J.M., Acosta, J.C., Jahn, G.A., Blanco, G., Laspiur, A., Ibagüengoytia, N.R., 2010. Reproductive biology of *Phymaturus cf. palluma*: a vulnerable lizard from the highlands of the Andes, San Juan, Argentina. *Herpetol. Conserv. Biol.* 5, 430–440.
- Cabrera, A.L., 1976. Enciclopedia Argentina de Agricultura y Jardinería, Tomo II, Fascículo 1: Regiones Fitogeográficas Argentinas. ACME, Buenos Aires.
- Cabrera, M.R., 2009. Lagartos del Centro de la Argentina. Edic. Indep., Córdoba.
- Cabrera, A.L., Willink, A., 1973. Biogeografía de América Latina. Monografía 13. Serie de Biología. OEA, Washington, DC.
- Cacciali, P., Scott, N.J., 2015. Key to the *Ophiodes* (Squamata: Sauria: Diploglossidae) of Paraguay with the description of a new species. *Zootaxa* 3980, 42–50.
- Carnevali, R., 1994. Fitoecografía de la Provincia de Corrientes. Cartas, escalas 1:500.000 y 1:1.000.000. Gobierno de la Provincia de Corrientes. INTA, Corrientes, Argentina.
- Cei, J.M., 1993. Reptiles del noroeste, nordeste y este de la Argentina. Herpetofauna de las selvas subtropicales, Puna y Pampas. Monografía XIV. Museo Regionale di Scienze Naturali, Torino, Italy.
- Colli, G.R., 1991. Reproductive ecology of *Ameiva ameiva* (Sauria, Teiidae), in the Cerrado of Central Brazil. *Copeia* 1991, 1002–1012.
- Crocco, M.C., Ibagüengoytia, N.R., Cussac, V., 2008. Contributions to the study of oviparity–viviparity transition: placental structures of *Liolaemus elongatus* (Squamata: Liolaemidae). *J. Morphol.* 269, 865–874.
- Derickson, W.K., 1976. Lipid storage and utilization in reptiles. *Am. Zool.* 16, 711–723.
- Doughty, P., Shine, R., 1998. Reproductive energy allocation and long-term energy stores in a viviparous lizard (*Eulamprus taylorianus*). *Ecology* 79, 1073–1083.
- Dunham, A.E., 1982. Demographic and life-history variation among populations of the iguanid lizard *Urosaurus ornatus*: implications for the study of life-history phenomena in lizards. *Herpetologica* 38, 208–221.
- Dunham, A.E., Miles, D.B., Reznick, D.N., 1988. Life history patterns in squamate reptiles. In: Gans, C., Huey, R.B. (Eds.), *Biology of the Reptilia*, vol. 16. Ecology B: Defense and Life History. Alan R. Liss, New York, pp. 441–522.
- Fitch, H.S., 1970. Reproductive cycles of lizards and snakes. *Univ. Kansas Mus. Nat. Hist. Misc. Publ.* 52, 1–247.
- Fitch, H.S., 1982. Reproductive cycles in tropical reptiles. *Occas. Pap. Mus. Nat. Hist. Univ. Kans.* 96, 1–53.
- Fitzgerald, L.A., Cruz, F.B., Perotti, G., 1999. Phenology of a lizard assemblage in the dry Chaco of Argentina. *J. Herpetol.* 33, 526–535.
- Goldberg, S.R., 1972. Seasonal weight and cytological changes in fat bodies and liver of the iguanid lizard *Sceloporus jarrovi* Cope. *Copeia* 1972, 227–232.
- Griffith, O.W., Blackburn, D.G., Brandley, M.C., Van Dyke, J.U., Whittington, C.M., Thompson, M.B., 2015. Ancestral state reconstructions require biological evidence to test evolutionary hypotheses: a case study examining the evolution of reproductive mode in squamate reptiles. *J. Exp. Zool. B (Mol. Dev. Evol.)* 324, 493–503.
- Guillette Jr., L.J., 1993. The evolution of viviparity in lizards. *Bioscience* 43, 742–751.
- Guillette Jr., L.J., Bearce, D.A., 1986. The reproductive and fat body cycles of the lizard, *Sceloporus grammicus disparilis*. *Trans. Kans. Acad. Sci.* 89, 31–39.
- Guillette Jr., L.J., Casas-Andreu, G., 1980. Fall reproductive activity in the high altitude Mexican lizard, *Sceloporus grammicus microlepidotus*. *J. Herpetol.* 14, 143–147.
- Guillette Jr., L.J., Casas-Andreu, G., 1981. Seasonal variation in fat body weights of the Mexican high elevation lizard *Sceloporus grammicus microlepidotus*. *J. Herpetol.* 15, 366–377.
- Guillette Jr., L.J., Casas-Andreu, G., 1987. The reproductive biology of the high elevation Mexican lizard *Barisia imbricata*. *Herpetologica* 43, 29–38.
- Guillette Jr., L.J., Méndez-de la Cruz, F.R., 1993. The reproductive cycle of the viviparous Mexican lizard *Sceloporus torquatus*. *J. Herpetol.* 27, 168–174.
- Hahn, W.E., Tinkle, D.W., 1965. Fat body cycling and experimental evidence for its adaptive significance to ovarian follicle development in the lizard *Uta stansburiana*. *J. Exp. Zool.* 158, 79–86.
- Herrel, A., Spithoven, L., Van Damme, R., De Vree, F., 1999. Sexual dimorphism of head size in *Gallotia galloti*: testing the niche divergence hypothesis by functional analyses. *Funct. Ecol.* 13, 289–297.
- Herrera, R., Voglio, D., Liotta, J., 2001. *Ophiodes intermedius* Boulenger, 1894 (Sauria: Anguillidae). *Cuad. Herpetol.* 15, 144.
- Hews, D.K., 1990. Examining hypotheses generated by field measures of sexual selection on male lizards, *Uta palmeri*. *Evolution* 44, 1956–1966.
- How, R.A., Shine, R., 1999. Ecological traits and conservation biology of five fossorial 'sand-swimming' snake species (*Simoselaps*: Elapidae) in south-western Australia. *J. Zool. Lond.* 249, 269–282.
- Hubert, J., 1985. The origin and development of oocytes. In: Gans, C. (Ed.), *Biology of the Reptilia*, vol. 14. Academic Press, New York, pp. 41–74.
- Ibagüengoytia, N.R., Cussac, V.E., 1998. Reproduction of the viviparous lizard *Liolaemus elongatus* in the highlands of southern South America: plastic cycles in response to climate? *Herpetol. J.* 8, 99–105.
- Ibagüengoytia, N.R., Cussac, V.E., 1999. Male response to low frequency of female reproduction in the viviparous lizard *Liolaemus* (Tropiduridae). *Herpetol. J.* 9, 111–117.
- Jackson, D.R., Franz, R., 1981. Ecology of the eastern coral snake (*Micrurus fulvius*) in northern peninsular Florida. *Herpetologica* 37, 213–228.
- James, C., Shine, R., 1985. The seasonal timing of reproduction: a tropical-temperate comparison in Australian lizards. *Oecologia* 67, 464–474.
- Janis, C.M., 1993. Tertiary mammal evolution in the context of changing climates, vegetation, and tectonic events. *Annu. Rev. Ecol. Syst.* 24, 467–500.
- Johnson, L.F., Jacob, J.S., Torrance, P., 1982. Annual testicular and androgenic cycles of the cottonmouth (*Agkistrodon piscivorus*) in Alabama. *Herpetologica* 38, 16–25.
- Jones, S.M., Wapstra, E., Swain, R., 1997. Asynchronous male and female gonadal cycles and plasma steroid concentrations in a viviparous lizard, *Niveoscincus ocellatus* (Scincidae), from Tasmania. *Gen. Comp. Endocrinol.* 108, 271–281.
- King, B., Lee, M.S.Y., 2015. Ancestral state reconstruction, rate heterogeneity, and the evolution of reptile viviparity. *Syst. Biol.* 64, 532–544.
- Leitão, M.D., 1973. Oviparidade em *Ophiodes striatus* (Spix, 1825), Sauria-Anguillidae. *Iheringia. Zool.* 42, 34–39.
- Leyton, V.C., Miranda, E.A., Bustos-Obregón, E., 1980. Gestational chronology in the viviparous lizard *Liolaemus gravenhorsti* (Gray) with remarks on ovarian and reproductive activity. *Arch. Biol.* 91, 347–361.
- Licht, P., 1971. Regulation of the annual testis cycle by photoperiod and temperature in the lizard *Anolis carolinensis*. *Ecology* 52, 240–252.
- Licht, P., 1973. Influence of temperature and photoperiod on the annual ovarian cycle in the lizard *Anolis carolinensis*. *Copeia* 1973, 465–472.
- Lozano, A., Uribe, M.C., Ramírez-Bautista, A., 2015. Seasonal and continuous spermatogenesis in the viviparous lizard *Sceloporus grammicus*: a study of two populations in contrasting environments from the Central Mexican Plateau. *Zool. Anz.* 254, 72–85.
- Marion, K.R., 1982. Reproductive cues for gonadal development in temperate reptiles: temperature and photoperiod effects on the testicular cycle of the lizard *Sceloporus undulatus*. *Herpetologica* 38, 26–39.
- Marques, O.A.V., Pizzatto, L., Almeida-Santos, S.M., 2013. Reproductive strategies of New World coral snakes, genus *Micrurus*. *Herpetologica* 69, 58–66.
- Mayhew, W.W., Wright, S., 1970. Seasonal changes in testicular histology of three species of the lizard genus *Uma*. *J. Morphol.* 130, 163–186.
- Medina, M., Ibagüengoytia, N.R., 2010. How do viviparous and oviparous lizards reproduce in Patagonia? A comparative study of three species of *Liolaemus*. *J. Arid Environ.* 74, 1024–1032.
- Méndez-de la Cruz, F.R., Villagrán, M., 1998. Reproducción asincrónica de *Sceloporus palaciosi* (Sauria: Phrynosomatidae) en México, con comentarios sobre sus ventajas y regulación. *Rev. Biol. Trop.* 46, 1159–1161.
- Méndez-de la Cruz, F.R., Villagrán-Santa Cruz, M., López-Ortiz, M.L., Hernández-Gallegos, O., 2013. Reproductive cycle of a high-elevation, oviparous lizard (*Sceloporus spinosus*: Reptilia: Phrynosomatidae). *Southwest. Nat.* 58, 54–63.
- Mesquita, D.O., Colli, G.R., 2010. Life history patterns in tropical South American lizards. In: Hernández-Gallegos, O., Méndez-de la Cruz, F.R., Méndez-Sánchez, J.F. (Eds.), *Reproducción en Reptiles: Morfología, Ecología y Evolución*. Universidad Autónoma del Estado de México, México, pp. 45–71.
- Miller, K.G., Wright, J.D., Fairbanks, R.G., 1991. Unlocking the ice house: Oligocene-Miocene oxygen isotopes, eustasy, and margin erosion. *J. Geophys. Res.* 96, 6829–6848.
- Mitchell, J.C., Zug, G.R., 1984. Spermatogenic cycle of *Nerodia taxipilota* (Serpentes: Colubridae) in south central Virginia. *Herpetologica* 40, 200–204.
- Mojica, B.H., Rey, B.H., Serrano, V.H., Ramírez-Pinilla, M.P., 2003. Annual reproductive activity of a population of *Cnemidophorus lemniscatus* (Squamata: Teiidae). *J. Herpetol.* 37, 35–42.
- Montechiaro, L., Kaffer, I.L., Quatros, F.C., Cechin, S., 2011. Feeding habits and reproductive biology of the glass lizard *Ophiodes cf. striatus* from subtropical Brazil. *North-West. J. Zool.* 7, 63–71.
- Mouton, P.F.N., Flemming, A.F., Stanley, E.L., 2012. Synchronized versus asynchronized breeding in cordylid lizards: an evolutionary perspective. *J. Zool.* 288, 191–198.
- Ortega-León, A.M., Villagrán-Santa Cruz, M., Zúñiga-Vega, J.J., Cueva del Castillo, R., Méndez-de la Cruz, F.R., 2009. Sperm viability in the reproductive tract of females in a population of *Sceloporus mucronatus* exhibiting asynchronous reproduction. *West. N. Am. Nat.* 69, 96–104.
- Ortiz, M.A., Boretto, J.M., Piantoni, C., Álvarez, B.B., Ibagüengoytia, N.R., 2014. Reproductive biology of the Amazon lava lizard (*Tropidurus torquatus*) from the Wet Chaco of Corrientes (Argentina): congeneric comparisons of ecotypic and inter-specific variations. *Can. J. Zool.* 92, 643–655.
- Ortiz, M.A., Boretto, J.M., Ibagüengoytia, N.R., 2016. Reproductive biology of the southernmost *Kentropyx* lizard from the Wet Chaco of Corrientes, Argentina. *Herpetol. J.* 26, 119–130.
- Ortiz-Jaureguizar, E., Cladera, G.A., 2006. Paleoenvironmental evolution of southern South America during the Cenozoic. *J. Arid Environ.* 66, 498–532.
- Pizzatto, L., 2005. Reproductive biology of the glass snake *Ophiodes fragilis* (Squamata: Anguillidae) in Southeast Brazil. *Herpetol. J.* 15, 9–13.
- Pough, F.H., Andrews, R.M., Crump, M.L., Savitzky, A.H., Wells, K.D., Brandley, M.C., 2016. *Herpetology*. Sinauer Associates, Sunderland, MA.
- Presst, M.R., 1994. Sexual size dimorphism and feeding energetics in *Anolis carolinensis*: why do females take smaller prey than males? *J. Herpetol.* 28, 292–298.
- Pyron, R.A., Burbrink, F.T., 2014. Early origin of viviparity and multiple reversions to oviparity in squamate reptiles. *Ecol. Lett.* 17, 13–21.
- Pyron, R.A., Burbrink, F.T., Wiens, J.J., 2013. A phylogeny and revised classification of Squamata, including 4161 species of lizards and snakes. *BMC Evol. Biol.* 13, 93.

- Quinn, H.R., 1979. Reproduction and growth of the Texas coral snake (*Micrurus fulvius tener*). *Copeia* 1979, 453–463.
- Ramírez-Bautista, A., Vitt, L.J., 1997. Reproduction in the lizard *Anolis nebulosus* (Polychrotidae) from the Pacific coast of Mexico. *Herpetologica* 53, 423–431.
- Ramírez-Bautista, A., Barba-Torres, J., Vitt, L.J., 1998. Reproductive cycle and brood size of *Eumeces lynx* from Pinal de Amoles, Queretaro, México. *J. Herpetol.* 32, 18–24.
- Ramírez-Bautista, A., Ramos-Flores, O., Sites Jr., J.W., 2002. Reproductive cycle of the spiny lizard *Sceloporus jarrovi* (Sauria: Phrynosomatidae) from north-central México. *J. Herpetol.* 36, 225–233.
- Ramírez-Bautista, A., Hernández-Ramos, D., Rojas-Martínez, A., Marshall, J.C., 2009. Fat bodies and liver mass cycles in *Sceloporus grammicus* (Squamata: Phrynosomatidae) from southern Hidalgo, México. *Herpetol. Conserv. Biol.* 4, 164–170.
- Ramírez-Bautista, A., Stephenson, B.P., Serrano Muñoz, C., Cruz-Elizalde, R., Hernández-Salinas, U., 2014. Reproduction and sexual dimorphism in two populations of the polymorphic spiny lizard *Sceloporus minor* from Hidalgo, México. *Acta Zool. (Stockholm)* 95, 397–408.
- Ramírez-Pinilla, M.P., 1991. Reproductive and fat body cycles of the viviparous lizard *Liolaemus huacahuasicus* (Sauria: Tropiduridae). *J. Herpetol.* 25, 205–208.
- Ramírez-Pinilla, M.P., 1992. Actividad reproductiva en tres especies simpátricas del género *Liolaemus* (Reptilia: Sauria: Tropiduridae). *Caldasia* 17, 67–74.
- Ramírez-Pinilla, M.P., 1995. Reproductive and fat body cycles of the oviparous lizard *Liolaemus bitaeniatus* (Sauria: Tropiduridae). *J. Herpetol.* 29, 256–260.
- Ramírez-Pinilla, M.P., Calderón-Espinosa, M.L., Flores-Villela, O., Muñoz-Alonso, A., Méndez de la Cruz, F.R., 2009. Reproductive activity of three sympatric viviparous lizards at Omiltemi Guerrero, Sierra Madre del Sur, México. *J. Herpetol.* 43, 409–420.
- Resende, F.C., Nascimento, L.B., 2015. The female reproductive cycle of the Neotropical snake *Atractus pantostictus* (Fernandes and Puerto, 1993) from south-eastern Brazil. *Anat. Histol. Embryol.* 44, 225–235.
- Retallack, G.J., 2001. Cenozoic expansion of grasslands and climatic cooling. *J. Geol.* 109, 407–426.
- Rocha, C.F.D., 1992. Reproductive and fat body cycles of the tropical sand lizard (*Liolaemus lutzae*) of southeastern Brazil. *J. Herpetol.* 26, 17–23.
- Rossetti, D.F., Bezerra, F.H.R., Dominguez, J.M.L., 2013. Late Oligocene-Miocene transgressions along the equatorial and eastern margins of Brazil. *Earth-Sci. Rev.* 123, 87–112.
- Schoener, T.W., 1967. The ecological significance of sexual dimorphism in size in the lizard *Anolis conspersus*. *Science* 155, 474–477.
- Schoener, T.W., Gorman, G.C., 1968. Some niche differences in three Lesser Antillean lizards of the genus *Anolis*. *Ecology* 49, 819–830.
- Sever, D.M., Hopkins, W.A., 2004. Oviductal sperm storage in the ground skink *Scincella laterale* Holbrook (Reptilia: Scincidae). *J. Exp. Zool.* 301A, 599–611.
- Shine, R., 1985. The evolution of viviparity in reptiles: an ecological analysis. In: Gans, C., Billett, F. (Eds.), *Biology of the Reptilia*. John Wiley & Sons, New York, pp. 605–694.
- Shine, R., 1988. The evolution of large body size in females: a critique of Darwin's fecundity advantage model. *Am. Nat.* 131, 124–131.
- Stewart, J.R., Castillo, R.E., 1984. Nutritional provision of the yolk of two species of viviparous reptiles. *Physiol. Zool.* 57, 377–383.
- Stewart, J.R., Thompson, M.B., 1993. A novel pattern of embryonic nutrition in a viviparous reptile. *J. Exp. Biol.* 174, 97–108.
- Stewart, J.R., Thompson, M.B., 2000. Evolution of placentation among squamate reptiles: recent research and future directions. *Comp. Biochem. Physiol. Part A* 127, 411–431.
- Strömberg, C.A.E., 2011. Evolution of grasses and grassland ecosystems. *Annu. Rev. Earth Planet. Sci.* 39, 517–544.
- Tinkle, D.W., Gibbons, J.W., 1977. The distribution and evolution of viviparity in reptiles. *Misc. Publ. Mus. Zool. Univ. Mich.* 154, 1–55.
- Uribe, M.C., 2002. Aparato reproductor masculino. In: Estrada-Flores, E., Uribe, M.C. (Eds.), *Atlas de Histología de Vertebrados*. Universidad Nacional Autónoma de México, México, D.F., pp. 133–148.
- Villagrán-Santa Cruz, M., Hernández-Gallegos, O., Méndez-de la Cruz, F.R., 2009. Reproductive cycle of the lizard *Sceloporus mucronatus* with comments on intraspecific geographic variation. *West. N. Am. Nat.* 69, 437–446.
- Vitt, L.J., 1983. Reproduction and sexual dimorphism in the tropical teiid lizard, *Cnemidophorus ocellifer*. *Copeia* 1983, 359–366.
- Vitt, L.J., 2015. Lizard reproduction: a history of discovery. In: Rheubert, J.L., Siegel, D.S., Trauth, S.E. (Eds.), *Reproductive Biology and Phylogeny of Lizards and Tuatara*. Taylor & Francis Group, Boca Raton, FL, pp. 1–26.
- Vitt, L.J., Caldwell, J.P., 2014. *Herpetology: An Introductory Biology of Amphibians and Reptiles*, 4th ed. Academic Press, San Diego.
- Vitt, L.J., Cooper Jr., W.E., 1985. The evolution of sexual dimorphism in the skink *Eumeces laticeps*: an example of sexual selection. *Can. J. Zool.* 63, 995–1002.
- Warne, R.W., Gilman, C.A., García, D.A., Wolf, B.O., 2012. Capital breeding and allocation to life-history demands are highly plastic in lizards. *Am. Nat.* 180, 130–141.
- Watling, J.L., Waddle, J.H., Kizirian, D., Donnelly, M.A., 2005. Reproductive phenology of three lizard species in Costa Rica, with comments on seasonal reproduction of Neotropical lizards. *J. Herpetol.* 39, 341–348.
- Webb, S.D., 1995. Biological implications of the Middle Miocene Amazon seaway. *Science* 269, 361–362.
- Wiens, J.J., Slingluff, J.L., 2001. How lizards turn into snakes: a phylogenetic analysis of body-form evolution in anguillid lizards. *Evolution* 55, 2303–2318.
- Zachos, J., Pagani, M., Sloan, L., Thomas, E., Billups, K., 2001. Trends, rhythms, and aberrations in global climate 65 Ma to present. *Science* 292, 686–693.